arXiv:math/0607141v3 [math.RA] 29 Apr 2008

(Co)Homology theories for oriented algebras

Juan Carlos Bustamante, Julie Dionne, and David Smith

Abstract. We study three different (co)homology theories for a family of pullbacks of algebras that we call oriented. We obtain a Mayer Vietoris long exact sequence of Hochschild and cyclic homology and cohomology groups for these algebras. We give examples showing that our sequence for Hochschild cohomology groups is different from the known ones. In case the algebras are given by quiver and relations, and that the simplicial homology and cohomology groups are defined, we obtain a similar result in a slightly wider context. Finally we also study the fundamental groups of the bound quivers involved in the pullbacks.

Introduction

Let k be a field and R be an associative k-algebra with unit. Given an $R-R$-bimodule ${}_RM_R$, or equivalently a right (or left) module over the enveloping algebra $R^e := R \otimes_k R^{op}$, the Hochschild homology and cohomology groups of R with coefficients in M_{R^e} are the groups $\mathrm{H}_n(R,M) = \mathrm{Tor}_n^{R^e}(M,R)$ and $\mathrm{H}^n(R,M) = \mathrm{Ext}^n_{R^e}(R,M)$, respectively. In case ${}_RM_R = {}_RR_R$ we simply write $\mathrm{H}_n(R)$, and $\mathrm{H}^n(R)$.

The low dimensional cohomological groups $\mathrm{H}^i(R)$ (that is $i \leq 2$) have clear classical interpretations in terms of the center, the derivations or the extensions of R. Besides this, these groups also play an important role in representation theory of algebras: the rigidity properties of R are closely related to $\mathrm{H}^2(R)$ and $\mathrm{H}^3(R)$, whereas $\mathrm{H}^1(R)$ gives information about the simple connectedness of R, and there are links between the vanishing of $\mathrm{H}^1(R)$ and $\mathrm{H}^2(R)$ and the representation type of R. Unfortunately, the groups $\mathrm{H}^i(R)$ are not easy to compute in general. There are some long exact sequences allowing to perform computations in some cases, see [**9, 23, 13, 14, 2**].

The main motivation for this work is to compute the Hochschild cohomology groups of the pullback R of two morphisms of k-algebras $A_1 \to C \leftarrow A_2$. However, as Examples 1.2.1 show, there is no easy immediate relation between the Hochschild cohomology groups of the algebras R, A_1, A_2 and C. Thus, we impose additional hypotheses on the morphisms above, and obtain a Mayer-Vietoris-like long exact sequence of Hochschild (co)homology groups of what we call **oriented algebras**. While doing so, one can see that using the same techniques it is possible to obtain

---

2000 *Mathematics Subject Classification.* 16E40.

Key words and phrases. Oriented algebras, pullbacks, Hochschild (co)homology, Cyclic (co)homology, Simplicial (co)homology, Fundamental group.

Corresponding author : D. Smith

analogous results about cyclic (co)homology as well as for simplicial (co)homology groups of algebras (if they are defined) and fundamental groups of the involved bound quivers, if the algebras are given in this way. This paper is organised as follows:

In Section 1 we recall some basic constructions related to bound quivers, and algebras given by the latter. We then define oriented algebras, give some examples, and make some preliminary observations.

Section 2 is devoted to Hochschild (co)homology. After recalling the basic definitions we establish the existence of two long exact sequences of Hochschild homology and cohomology groups, in Theorems 2.2.3 and 2.3.2. Moreover, we show that the morphisms involved in the cohomological sequence are compatible with the Gerstenhaber algebra structure of $\mathrm{H}^*(R) := \bigoplus_{i\geq 0}\mathrm{H}^i(R)$. The remaining part of this section is devoted to specialize our results to the case where the algebra C is what we call a core algebra, and, on the other hand, to compare the long exact sequence of Theorem 2.3.2 with the known long exact sequences of Hochschild cohomology groups of [**9, 23, 13, 14, 2**].

In Section 3, again, after recalling some constructions, we establish the existence of long exact sequences of cyclic homology and cohomology groups, in Theorems 3.3.5 and 3.3.8. Moreover, we show that these sequences are compatible with Connes' long exact sequences relating Hochschild and cyclic (co)homology groups.

Finally, in Section 4 we turn our attention to algebras given by bound quivers, and, more precisely to algebras having semi-normed basis. The existence of such basis allows to define the simplicial (co)homology groups of algebras, following [**21**]. In the same flavor of the previous parts of this paper, but with weaker hypotheses, we establish the existence of long exact sequences of simplicial (co)homology groups, in Propositions 4.1.1 and 4.1.2. We then consider fundamental groups of bound quivers, and obtain an explicit formula that allows to compute them in a context which is slightly different from that of the oriented pullbacks.

1. Preliminaries

1.1. Algebras and quivers. Throughout this paper, k denotes a commutative field and all algebras are associative k-algebras with unit. Tensor products are taken over k unless otherwise stated, so $\otimes = \otimes_k$. Let R^{op} be the opposite algebra of R, and denote by R^e its enveloping algebra $R\otimes R^{op}$ of R. We shall freely use the fact that the category of R-bimodules is equivalent to the category of right (or left) R^e-modules: given a R-bimodule M, define the right R^e-structure by $m(a\otimes b) = bma$ (and the left R^e-structure by $(a\otimes b)m = amb$).

A (finite) **quiver** is a quadruple $Q = (Q_0, Q_1, s, t)$, where Q_0 and Q_1 are two (finite) sets, respectively called the **set of vertices** and the **set of arrows**, and s and t are two maps from Q_1 to Q_0 associating to each arrow its **source** and its **target**, respectively. A quiver Q is **connected** if its underlying graph is connected.

A **path** on **length** n is a sequence of n arrows $\omega = \alpha_1\alpha_2\cdots\alpha_n$ such that $t(\alpha_i) = s(\alpha_{i+1})$ for each i, with $1 \leq i \leq n-1$. The **source** of such a path is $s(\alpha_1)$, and its **target** is $t(\alpha_n)$. Also, to each vertex x of Q, we associate the **stationary path** ε_x, whose length is 0, and whose source and target are x. An **oriented cycle** is a path $\omega = \alpha_1\alpha_2\cdots\alpha_n$ of length at least one, such that $s(\alpha_1) = t(\alpha_n)$.

The **path algebra** kQ is the k-vector space having as basis the set of all paths in Q endowed with the following multiplication: let ω_1 and ω_2 be two basis elements, then $\omega_1\omega_2$ is their composition if $t(\omega_1) = s(\omega_2)$, and 0 otherwise. A **relation** from x to y, with $x, y \in Q_0$, is a linear combination $\rho = \sum_{i=1}^{r} \lambda_i \omega_i$ where, for each i, ω_i is a path of length at least two from x to y in Q and $\lambda_i \in k \setminus \{0\}$. Let F be the two-sided ideal of kQ generated by all the arrows of Q. An ideal I of kQ is **admissible** if there exists $m \geq 2$ such that $F^m \subseteq I \subseteq F^2$. In this case, the pair (Q, I) is called a **bound quiver**. The algebra kQ/I is basic, connected (if so is Q), and finite dimensional (if Q is finite). By abuse of notation, we simply denote the primitive pairwise orthogonal idempotents $\varepsilon_x + I$ of kQ/I by ε_x, so that $1_{kQ/I} = \sum_{x \in Q_0} \varepsilon_x$.

Conversely, for every finite dimensional, connected and basic algebra A over an algebraically closed field k, there exists a unique connected quiver Q and surjective maps $\nu : kQ \to A$ with $I = \operatorname{Ker} \nu$ admissible (see [**4**]). The pair (Q, I) is called a **presentation** of A. Following [**4**], any basic algebra $A = kQ/I$ can equivalently be regarded as a locally bounded k-category having as object class the set $A_0 = Q_0$ and as morphism set from x to y the k-vector space $A(x, y) = \varepsilon_x A \varepsilon_y$. In what follows, we use both terminologies, and pass from one to the other freely. A subcategory B of A is called **full** if, for all $x, y \in B_0$, we have $B(x, y) = A(x, y)$, and **convex** if any vertex which lies on a path in Q having its source and target in B_0 also belongs to B_0. Note that convexity does not imply fullness.

1.2. Pullbacks. Let $f_1 : A_1 \to C$ and $f_2 : A_2 \to C$ be two surjective morphisms of k-algebras. Then the set $R = \{(a_1, a_2) \in A_1 \times A_2 | \ f_1(a_1) = f_2(a_2)\}$, endowed with the natural operations becomes a k-algebra, which is in fact the pullback of f_1 and f_2. As Examples 1.2.1 below show, one cannot hope to have a direct relation between the Hochschild cohomology groups of R, A_1, A_2, and C in general.

In [**17**], a special kind of pullbacks has been considered: assume that $1_{A_1} = u_1' + u_1''$, $1_{A_2} = u_2' + u_2''$ and that there exists an isomorphism $A_1/ < u_1' > \simeq A_2/ < u_2' >$ sending the image of u_1'' under the projection, to the image of u_2''. Let C be this common quotient, f_1 and f_2 the projections, and, as above, R be the pullback of f_1 and f_2. In R, let $e_1' = (u_1', 0)$, $e_2' = (0, u_2')$, and $e = (u_1'', u_2'')$. A direct computation shows that $e_1' R e_2' = e_2' R e_1' = 0$ and that $1_R = e_1' + e + e_2'$. In [**17**] one can find the detailed construction of the quiver of R from those of A_1, A_2 and C.

Examples 1.2.1.

(a) Let A_1 and $A_2 = A_1^{op}$ be the hereditary algebras given by the quivers $1 \xrightarrow{\alpha} 2$ and $1 \xleftarrow{\beta} 2$, respectively. Moreover, let C be the semisimple algebra generated by the vertices 1 and 2, and, for $i \in \{1, 2\}$ let $f_i : A_i \to C$ be the projections. The pullback R is then the bound quiver algebra given by the quiver $1 \underset{\beta}{\overset{\alpha}{\rightleftarrows}} 2$ with relations $\alpha\beta = 0 = \beta\alpha$. For each $n > 0$, we have $\mathrm{H}^n(R) \neq 0$ by [**8**, (p. 96)], while $\mathrm{H}^n(A_1) = \mathrm{H}^n(A_2) = \mathrm{H}^n(C) = 0$.

(b) Let R be the algebra given by the quiver

$$\begin{tikzcd}
 & 2 \arrow[dr, "\beta"] & \\
1 \arrow[ur, "\alpha"] & & 3 \arrow[dl, "\gamma"] \\
 & 4 \arrow[ul, "\delta"] &
\end{tikzcd}$$

bounded by the ideal generated by all paths of length two. Let A_1 and A_2 be full subcategories of R generated by the sets of vertices $\{1,\ 2,\ 4\}$, and $\{2,\ 3,\ 4\}$ respectively. Then, R is the pullback of the projections of A_1 and A_2 over the semisimple algebra C generated by the vertices 1 and 3. By setting $e = \varepsilon_2 + \varepsilon_4$, $e'_1 = \varepsilon_1$ and $e'_2 = \varepsilon_3$, it is easily seen that $1_R = e'_1 + e + e'_2$ and $e'_1Re'_2 = 0 = e'_2Re'_1$.
Again, as in the previous example, we get from [**8**, (p. 96)] that $\mathrm{H}^n(R) \neq 0$ for infinitely many $n > 0$, while $\mathrm{H}^n(A_1) = \mathrm{H}^n(A_2) = \mathrm{H}^n(C) = 0$ for each $n > 0$. In fact what allows to construct nontrivial elements in $\mathrm{H}^{4n}(R)$ is the fact that $eRe'_i \neq 0$ and $e'_iRe \neq 0$ for $i \in \{1, 2\}$.

The last example motivates the following definition.

DEFINITION 1.2.2. *Let R be an algebra. Assume that there exists a decomposition $1_R = e'_1 + e + e'_2$ of the unit of R, where e, e'_1, e'_2 are orthogonal idempotents of R. Then R is said to be* **oriented** *by e, e'_1, e'_2 if*

$$e'_1Re'_2 = e'_2Re'_1 = 0 \tag{1.1}$$

and one of the following conditions is satisfied :

$$eRe'_1 = eRe'_2 = 0; \tag{1.2}$$

$$e'_1Re = e'_2Re = 0; \tag{1.3}$$

$$e'_1Re = eRe'_2 = 0. \tag{1.4}$$

In this case, we set $e_i = e'_i + e, A'_i = e'_iRe'_i$, $A_i = e_iRe_i$ for $i = 1, 2$ and $C = eRe$. Finally, we set $E = ke'_1 + ke + ke'_2 \subseteq R$, $E_1 = ke'_1 + ke \subseteq A_1$, $E_2 = ke'_2 + ke \subseteq A_2$ and $E_C = ke \subseteq C$.

Clearly, R, C, A_1 and A_2 are E-bimodules, and C, A_1 and A_2 are also bimodules over the semisimple algebras E, E_1 and E_2 respectively. As we shall see, these semisimple algebras will play a crucial role in the sequel, as nice substitutes for k.

Outside the context of bound quivers, an algebra Λ is said to be **convex** in bigger algebra Γ ($\Lambda \subseteq \Gamma$, but not necessarily a subalgebra) if whenever $\lambda_1, \lambda_2, \dots, \lambda_n$ are elements in Γ such that $1_\Lambda\lambda_1\lambda_2\cdots\lambda_n1_\Lambda \neq 0$, then each of the λ_i's belong to Λ.

Also observe that conditions (1.1)-(1.4) are chosen in such a way that if R is oriented by e, e'_1, e'_2, then the algebras A_i, A'_i and C are convex in R, for $i = 1, 2$.

EXAMPLES 1.2.3.

(a) If $e = 0$ but $e'_1 \neq 0$ and $e'_2 \neq 0$, then condition (1.1) simply means that $R \simeq A'_1 \times A'_2 = A_1 \times A_2$.

(b) Let Γ, Λ be two k-algebras and ${}_\Gamma M_\Lambda$ be a $\Gamma - \Lambda$-bimodule. We can then construct the **triangular matrix ring** $R = \begin{pmatrix} \Gamma & 0 \\ {}_\Lambda M_\Gamma & \Lambda \end{pmatrix}$. By taking $e'_1 = \begin{pmatrix} 1_\Gamma & 0 \\ 0 & 0 \end{pmatrix}$, $e'_2 = 0$ and $e = \begin{pmatrix} 0 & 0 \\ 0 & 1_\Lambda \end{pmatrix}$ one can see that R is oriented. In case Λ is a division ring, the R is called a **one-point extension** of Γ by M. See Remark 1.2.4, (a), case 1.3.

REMARK 1.2.4.

(a) If R is oriented by e, e'_1 and e'_2, using the decomposition $1_R = e'_1 + e + e'_2$, the relation $R = 1_R \cdot R \cdot 1_R$, and the isomorphism of k-algebras $R \simeq \mathrm{End}_R(R)$, one can write R in a matrix form. The next figure shows the possible matrix configurations, together with their corresponding quiver configurations in case R is a bound quiver algebra. Note that the only possible nonzero paths in each case are indicated by the depicted arrows.

$$\begin{pmatrix} A_1' & e_1'Re & 0 \\ 0 & C & 0 \\ 0 & e_2'Re & A_2' \end{pmatrix} \qquad \begin{pmatrix} A_1' & 0 & 0 \\ eRe_1' & C & eRe_2' \\ 0 & 0 & A_2' \end{pmatrix} \qquad \begin{pmatrix} A_1' & 0 & 0 \\ eRe_1' & C & 0 \\ 0 & e_2'Re & A_2' \end{pmatrix}$$



case (1.2)



case (1.3)



case (1.4)

(b) Note that any matrix algebra of one of these forms is oriented by the idempotents $e_1' = e_{11}, e = e_{22}, e_2' = e_{33}$, where e_{ii} is the 3×3-matrix having a 1 in position (i,i) and 0 elsewhere. Moreover, observe that in this case R is the pullback of the projections $\pi_i : A_i \to C : e_i r e_i \mapsto ere$, for $i = 1, 2$. Therefore, the oriented algebras (with e, e_1' and e_2' nonzero) generalize the "Nakayama oriented pullbacks" introduced in [**18**]. However, not all pullbacks are oriented, as Examples 1.2.1 show.

1.3. Some immediate observations. In this section, we establish some immediate results that will be needed in the sequel. From now on we adopt the notations and terminology of Definition 1.2.2.

REMARK 1.3.1. Let R be an algebra oriented by a set of idempotents e, e_1', e_2'.

(a) Let $\sigma_1, \sigma_2, \tau_1, \tau_2 \in \{e_1', e, e_2'\}$. Then, in $R \otimes_E R$, we have

$$\sigma_1 R \tau_1 \otimes_E \sigma_2 R \tau_2 = \sigma_1 R \tau_1 \sigma_2 \otimes_E \sigma_2 R \tau_2 = \sigma_1 R \tau_1 \otimes_E \tau_1 \sigma_2 R \tau_2$$

and, since the idempotents e_1', e and e_2' are orthogonal, this vanishes if $\tau_1 \neq \sigma_2$. The same argument shows that $R^{\otimes_E n}$ is generated by the elements of the form $r_1 \otimes r_2 \otimes \cdots \otimes r_n$, with $r_i \in \sigma_i R \tau_i$ with $\sigma_i, \tau_i \in \{e, e_1', e_2'\}$ and $\tau_i = \sigma_{i+1}$ for each i. Note that we are strongly using the fact that the tensor products are taken over E. The claim does not hold if tensors are taken over k.

(b) As noted before, C is full and convex in A_1 and A_2, which in turn are full and convex in R. Thus, we have fully faithful embeddings $\mathrm{mod} C \subseteq \mathrm{mod} A_i \subseteq \mathrm{mod} R$. Consequently, $X \otimes_E Y \simeq X \otimes_{E_C} Y$ whenever X and Y are modules over C (or C^{op}).

(c) Let X be an R-bimodule, or, equivalently an R^e-module. Then there exists an epimorphism $p : (R^e)^n \to X$, so that for $x \in X$ we have $x = p(r)$, hence

$$e_1' x e_2' = (e_1' \otimes e_2') x = (e_1' \otimes e_2') p(r) = p((e_1' \otimes e_2') r) = p(e_1' r e_2') = p(0) = 0$$

This shows that $e_1' X e_2' = 0$. Similarly, one can show that $e_2' X e_1' = 0$.

We then get the following lemmata

LEMMA 1.3.2. *If R is oriented by a set of idempotents e, e_1' and e_2', then*

(a) $eR^{\otimes_E n} e = (eRe)^{\otimes_E n} = C^{\otimes_E n}$;

(b) $e_1 R^{\otimes_E n} e_1 = (e_1 R e_1)^{\otimes_E n} = A_1^{\otimes_E n}$;

(c) *$e_2R^{\otimes_E n}e_2 = (e_2Re_2)^{\otimes_E n} = A_2^{\otimes_E n}$;*

PROOF. We only prove (a) since the proofs of (b) and (c) are similar.
(a). Clearly, eRe is included in R, whence $(eRe)^{\otimes_E n} \subseteq R^{\otimes_E n}$. Multiplying by e on both sides gives $(eRe)^{\otimes_E n} \subseteq eR^{\otimes_E n}e$. On the other hand, let $\tau = e(r_1 \otimes r_2 \otimes \cdots \otimes r_n)e$ be a nonzero element of $eR^{\otimes_E n}e$. It then follows from Remark 1.3.1 (a) and the convexity of C in R that $r_i \in eRe$, for each i. Thus $\tau \in (eRe)^{\otimes_E n}$. □

LEMMA 1.3.3. *If R is oriented by a set of idempotents e, e'_1 and e'_2, then the following hold, for $i = 1, 2$:*

(a) *If $eRe'_i = 0$, then $A_i^{\otimes_E n} = e'_iR^{\otimes_E n}e'_i \oplus e'_iR^{\otimes_E n}e \oplus C^{\otimes_E n}$;*
(b) *If $e'_iRe = 0$, then $A_i^{\otimes_E n} = e'_iR^{\otimes_E n}e'_i \oplus eR^{\otimes_E n}e'_i \oplus C^{\otimes_E n}$.*

PROOF. This follows from $A_i^{\otimes_E n} = (e_iRe_i)^{\otimes_E n} = e_iR^{\otimes_E n}e_i = (e'_i+e)R^{\otimes_E n}(e'_i+e)$ and Remark 1.3.1 (c). □

2. Hochschild (co)homology

2.1. Definitions and notations. Let, as before, k be a field and R be a k-algebra. For each $n \geq 2$, consider the morphism of R-bimodules $b' : R^{\otimes n+1} \to R^{\otimes n}$ given by

$$b'(r_0 \otimes \cdots \otimes r_n) \quad := \quad \sum_{i=0}^{n-1}(-1)^i(r_0 \otimes \cdots \otimes r_ir_{i+1} \otimes \cdots \otimes r_n) \tag{2.1}$$

Since $b' \circ b' = 0$, we get a complex $(R^{\otimes\bullet}, b')$. The maps $h : R^{\otimes n} \to R^{\otimes n+1}$ given by $h(r_0 \otimes \cdots \otimes r_{n-1}) = 1 \otimes r_0 \otimes \cdots \otimes r_{n-1}$ define a contracting homotopy, so $(R^{\otimes\bullet}, b')$ is a resolution of the R^e-module R, the so-called **Hochschild resolution**.

Let M be a R-bimodule. If we apply the functor $M \otimes_{R^e} -$ to the Hochschild resolution of R, we get a complex whose nth homology group is, by definition, the nth **Hochschild homology group** of R with coefficients in M, denoted $\mathrm{H}_n(R, M)$. Similarly, the nth **Hochschild cohomology group** $\mathrm{H}^n(R, M)$ is the nth homology group of the complex obtained upon applying the functor $\mathrm{Hom}_{R^e}(-, M)$ to the Hochschild resolution of R. We simply write $\mathrm{H}_n(R)$ and $\mathrm{H}^n(R)$ in case $M = R$.

Since k is a field, $R^{\otimes n-2}$ is k-projective and so the module $R^{\otimes n}$ is R^e-projective for $n \geq 2$. Then the Hochschild resolution $(R^{\otimes\bullet}, b')$ is a projective resolution of R over R^e. So in our context one may define the Hochschild (co)homology groups of R with coefficients in M in the following way:

$$\mathrm{H}_n(R, M) = \mathrm{Tor}_n^{R^e}(M, R) \qquad \text{and} \qquad \mathrm{H}^n(R, M) = \mathrm{Ext}^n_{R^e}(R, M)$$

Thus, for each $n \geq 0$ there exists an isomorphism $\phi_n : D\mathrm{H}^n(R, DM) \simeq \mathrm{H}_n(R, M)$ (see [**6**, p. 181]), where $D = \mathrm{Hom}_k(-, k)$ is the usual duality. Moreover, the Hochschild (co)homology groups do not depend on the projective resolution we consider to compute them. For our purpose, it will be necessary to deal with a different projective resolution of ${}_RR_R$. In particular we will use the following lemma, from [**24**].

LEMMA 2.1.1. *Let R be an algebra and E be a semisimple subalgebra of R. The complex*

$$(R^{\otimes_E\bullet}, b') : \quad \cdots \longrightarrow R^{\otimes_E n+1} \xrightarrow{b'} R^{\otimes_E n} \xrightarrow{b'} \cdots \xrightarrow{b'} R^{\otimes_E 2} \xrightarrow{b'} R \longrightarrow 0 \tag{2.2}$$

where b' is given by the formula (2.1) in which we replace the tensor products over k by tensor products over E, is a projective resolution of R as R^e-module.

Hence the groups $\mathrm{H}_n(R,M)$ (or $\mathrm{H}^n(R,M)$) are the homology groups of the complex obtained when we apply the functor $M\otimes_{R^e}-$ (or $\mathrm{Hom}_{R^e}(-,M)$, respectively) to the projective resolution (2.2).

Now, for $n\geq 2$, we have canonical isomorphisms

$$\begin{array}{rcl} M\otimes_{R^e}R\otimes_E R^{\otimes_E n-2}\otimes_E R & \simeq & M\otimes_{E^e}R^{\otimes_E n-2} \\ m\otimes r\otimes x\otimes r' & \mapsto & r'mr\otimes x \end{array} \tag{2.3}$$

and

$$\begin{array}{rcl} \mathrm{Hom}_{R^e}(R\otimes_E R^{\otimes_E n-2}\otimes_E R,M) & \simeq & \mathrm{Hom}_{E^e}(R^{\otimes_E n-2},M) \\ f & \mapsto & \overline{f}:m\mapsto f(1_R\otimes m\otimes 1_R) \end{array} \tag{2.4}$$

Using these identifications we obtain the following (compare with [**7**, (1.2)]):

Proposition 2.1.2. *Let R be an algebra and E be a semisimple subalgebra of R. Let M be a R-bimodule.*

(a) *The Hochschild homology groups $\mathrm{H}_n(R,M)$ are the homology groups of the complex*

$$M\otimes_{E^e}R^{\otimes_E\bullet}\ :\ \cdots\to M\otimes_{E^e}R^{\otimes_E 3}\xrightarrow{\delta_2}M\otimes_{E^e}R^{\otimes_E 2}\xrightarrow{\delta_1}M\otimes_{E^e}R\xrightarrow{\delta_0}M\otimes_{E^e}E\to 0$$

where the differential $\delta_\bullet$ is deduced from the definition of b' (2.2), and the formula (2.3).

(b) *The Hochschild cohomology groups $\mathrm{H}^n(R,M)$ are the homology groups of the complex*

$$\mathrm{Hom}_{E^e}(R^{\otimes\bullet},M)\ :\ 0\to\mathrm{Hom}_{E^e}(E,M)\xrightarrow{\delta^0}\mathrm{Hom}_{E^e}(R,M)\xrightarrow{\delta^1}\mathrm{Hom}_{E^e}(R^{\otimes_E 2},M)\xrightarrow{\delta^2}\cdots$$

where, the differential $\delta^\bullet$ is deduced from the definition of b' (2.2), and the formula (2.4).

The Yoneda product in $\mathrm{H}^*(R)=\bigoplus_{i\geq 0}\mathrm{H}^i(R)$ is induced by a cup-product defined at the cochain level in the complex of Proposition 2.1.2 (b), as follows. Given $f\in\mathrm{Hom}_{E^e}(R^{\otimes_E n},R)$ and $g\in\mathrm{Hom}_{E^e}(R^{\otimes_E m},R)$ we define the element $f\cup g$ in $\mathrm{Hom}_{E^e}(R^{\otimes_E n+m},R)$ by the rule:

$$(f\cup g)(r_1\otimes\cdots\otimes r_{n+m})=f(r_1\otimes\cdots\otimes r_n)g(r_{n+1}\otimes\cdots\otimes r_{n+m}).$$

On the other hand, given $i\in\{1,\ldots,n\}$, define $f\circ_i g\in\mathrm{Hom}_{E^e}(R^{\otimes_E n+m-1},R)$ by the rule:

$$(f\circ_i g)(r_1\otimes\cdots\otimes r_{n+m-1})=f(r_1\otimes\cdots\otimes g(r_i\otimes\cdots\otimes r_{i+m-1})\otimes\cdots\otimes r_{n+m-1}).$$

Then we define the composition

$$f\circ g=\begin{cases}0, & \text{if } n=0;\\ \sum_{i=1}^{n}(-1)^{(i-1)(m-1)}f\circ_i g, & \text{if } n>0.\end{cases}$$

and finally the bracket of f and g as $[f,g]=f\circ g-(-1)^{(n-1)(m-1)}g\circ f$. As the cup-product, this bracket induces an operation at the cohomology level. These two structures are related: in fact $\mathrm{H}^*(R)=\bigoplus_{i\geq 0}\mathrm{H}^i(R)$, refered as the **Hochschild algebra** over R in the sequel, is a Gerstenhaber algebra (see [**12**]). That is $\mathrm{H}^*(R)$ is a graded k-vector space endowed with a product $\cup$ which makes it a graded commutative algebra, and a Lie bracket $[-,-]$ of degree -1 that makes of $\mathrm{H}^*(R)$ a graded Lie algebra, and such that $[x, y \cup z] = [x,y] \cup z + (-1)^{(|x|-1)|y|} y \cup [x,z]$, where $|x|$ and $|y|$ are the degrees of x and y, respectively.

2.2. Long exact sequence of Hochschild homology groups. Let R be an algebra, oriented by a set of idempotents e, e_1' and e_2' (see Definition 1.2.2). In this section, we prove our first main result, that is the existence of a long exact sequence relating the Hochschild homology groups of R, C, A_1 and A_2. This will be a consequence of the following proposition, in which we freely use the identifications given in Remark 1.3.1 and Lemma 1.3.2.

PROPOSITION 2.2.1. *Let R be an algebra, oriented by three idempotents e, e_1' and e_2'. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_2 = e_2Re_2$ be as in Definition 1.2.2. Let M be an R-bimodule. For any $n \geq 0$, the sequence*

$$0 \longrightarrow eMe \bigotimes_{E_C^e} C^{\otimes_{E_C} n} \xrightarrow{\alpha^n} \bigoplus_{i=1}^{2} e_iMe_i \otimes_{E_i^e} A_i^{\otimes_{E_i} n} \xrightarrow{\beta^n} M \bigotimes_{E^e} R^{\otimes_E n} \longrightarrow 0$$

is exact in $\mathrm{mod}E^e$, *where $\alpha^n = (\alpha_1^n, \alpha_2^n)^t$, with $\alpha_i^n(eme \otimes exe) = e_iemee_i \otimes e_iexee_i$, and $\beta^n = (\beta_1^n, \beta_2^n)$, with $\beta_i^n(e_ime_i \otimes e_ixe_i) = (-1)^i(e_ime_i \otimes e_ixe_i)$, for $i = 1, 2$.*

PROOF. Since R is oriented, we have three cases to consider, these are (1.2)-(1.4). We only prove the exactness for the case (1.2) since the proofs for the other cases are similar. So, assume that $e_1'Re_2' = e_2'Re_1' = 0$ and also $eRe_1' = eRe_2' = 0$. Using Remark 1.3.1 (c), we have $e_1'Xe_2' = e_2'Xe_1' = 0$, and similarly $e_1'Xe = e_2'Xe = 0$ for any R-bimodule X.

Using $1_R = e_1' + e + e_2'$, we have the decomposition :

$$R^{\otimes_E n} = e_1'R^{\otimes_E n}e_1' \oplus eR^{\otimes_E n}e_1' \oplus eR^{\otimes_E n}e \oplus eR^{\otimes_E n}e_2' \oplus e_2'R^{\otimes_E n}e_2'$$

and so

$$\begin{aligned}
M \otimes_{E^e} R^{\otimes_E n} &= (M \otimes_{E^e} e_1'R^{\otimes_E n}e_1') \oplus (M \otimes_{E^e} eR^{\otimes_E n}e_1') \\
&\quad \oplus (M \otimes_{E^e} eR^{\otimes_E n}e) \\
&\quad \oplus (M \otimes_{E^e} eR^{\otimes_E n}e_2') \oplus (M \otimes_{E^e} e_2'R^{\otimes_E n}e_2') \\
&\simeq (e_1Me_1 \otimes_{E_1^e} e_1'R^{\otimes_E n}e_1') \oplus (e_1Me_1 \otimes_{E_1^e} eR^{\otimes_E n}e_1') \\
&\quad \oplus (eMe \otimes_{E_C^e} C^{\otimes_{E_C} n}) \\
&\quad \oplus (e_2Me_2 \otimes_{E_2^e} eR^{\otimes_E n}e_2') \oplus (e_2Me_2 \otimes_{E_2^e} e_2'R^{\otimes_E n}e_2')
\end{aligned}$$

where the isomorphism follows from immediate identification.

Similarly, for $i = 1, 2$, we get from Lemma 1.3.3, that

$$\begin{aligned}
e_iMe_i \otimes_{E_i^e} A_i^{\otimes_{E_i} n} &\simeq (e_iMe_i \otimes_{E_i^e} e_i'R^{\otimes_E n}e_i') \oplus (e_iMe_i \otimes_{E_i^e} eR^{\otimes_E n}e_i') \\
&\quad \oplus (eMe \otimes_{E_C^e} C^{\otimes_{E_C} n})
\end{aligned}$$

With these isomorphisms and identifications in mind, we obtain the desired short exact sequence of complexes. □

REMARK 2.2.2. In the above proposition, the fact that the tensor products are taken over E is crucial. We have used Lemmata 1.3.2 and 1.3.3, which do not hold true if tensor products are taken over k.

We can know prove our first main result.

Theorem 2.2.3. *Let R be an algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_1 = e_2Re_2$ be as in Definition 1.2.2. For any R-bimodule M, there exists a long exact sequence of Hochschild homology groups :*

$$\cdots \longrightarrow \mathrm{H}_1(C, eMe) \longrightarrow \mathrm{H}_1(A_1, e_1Me_1) \oplus \mathrm{H}_1(A_2, e_2Me_2) \longrightarrow \mathrm{H}_1(R, M) \longrightarrow$$
$$\longrightarrow \mathrm{H}_0(C, eMe) \longrightarrow \mathrm{H}_0(A_1, e_1Me_1) \oplus \mathrm{H}_0(A_2, e_2Me_2) \longrightarrow \mathrm{H}_0(R, M) \longrightarrow 0$$

Proof. The result follows from the above Proposition and the observation that α^n and β^n commute with the differential δ_n of Proposition 2.1.2 (a), giving rise to a short exact sequence of complexes.

$$0 \longrightarrow eMe \otimes_{E^e_C} C^{\otimes_{E_C}\bullet} \xrightarrow{\alpha^\bullet} \bigoplus_{i=1}^{2} e_iMe_i \otimes_{E^e_i} A_i^{\otimes_{E_i}\bullet} \xrightarrow{\beta^\bullet} M \otimes_{E^e} R^{\otimes_E \bullet} \longrightarrow 0$$

whose homology groups are the desired Hochschild homology groups. □

2.3. Long exact sequence of Hochschild cohomology groups. In this section, we show an analogue to Theorem 2.2.3 for Hochschild cohomology groups. In addition, we show that the maps involved in the long exact sequence are compatible with the Yoneda product and the Lie bracket of the involved Hochschild algebras.

At this point, we stress that one can deduce a long exact sequence of Hochschild cohomology groups by applying the isomorphism $\phi_n : D\mathrm{H}^n(R, DM) \simeq \mathrm{H}_n(R, M)$ on the long exact sequence of Theorem 2.2.3. We will however avoid this approach since we need a clear description of the morphisms to verify their compatibility with the Yoneda products and Lie brackets. Their knowledge will also be important for Theorem 3.3.9. Nevertheless, since our approach is similar to the one we have used for Hochschild homology, we skip some details, for instance the proof of the following proposition, which is an analogue to Proposition 2.2.1.

Proposition 2.3.1. *Let R be an algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_1 = e_2Re_2$ be as in Definition 1.2.2. Let M be an R-bimodule. For any $n \geq 0$, the sequence*

$$0 \to \mathrm{Hom}_{E^e}(R^{\otimes_E n}, M) \overset{\alpha^n}{\to} \bigoplus_{i=1}^{2} \mathrm{Hom}_{E^e_i}(A_i^{\otimes_{E_i} n}, e_iMe_i) \overset{\beta^n}{\to} \mathrm{Hom}_{E^e_C}(C^{\otimes_{E_C} n}, eMe) \to 0$$

is exact in $\mathrm{mod}E^e$, *where* $\alpha^n = (\alpha^n_1, \alpha^n_2)^t$, *with* $\alpha^n_i(f)(r_1 \otimes \cdots \otimes r_n) = f(e_ir_1 \otimes \cdots \otimes r_ne_i)$ *and* $\beta^n = (\beta^n_1, \beta^n_2)$, *with* $\beta^n_i(f)(r_1 \otimes \cdots \otimes r_n) = (-1)^i f(er_1 \otimes \cdots \otimes r_ne)$, *for* $i = 1, 2$.

We can now prove our second main result.

Theorem 2.3.2. *Let R be an algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_1 = e_2Re_2$ be as in Definition 1.2.2. For any R-bimodule M, there exists a long exact sequence of Hochschild cohomology groups:*

$$0 \longrightarrow \mathrm{H}^0(R, M) \xrightarrow{\alpha^0} \mathrm{H}^0(A_1, e_1Me_1) \oplus \mathrm{H}^0(A_2, e_2Me_2) \xrightarrow{\beta^0} \mathrm{H}^0(C, eMe) \longrightarrow$$
$$\longrightarrow \mathrm{H}^1(R, M) \xrightarrow{\alpha^1} \mathrm{H}^1(A_1, e_1Me_1) \oplus \mathrm{H}^1(A_2, e_2Me_2) \xrightarrow{\beta^1} \mathrm{H}^1(C, eMe) \longrightarrow \cdots$$

In addition, if $M = R$, then the induced maps $\alpha_i^n : \mathrm{H}^n(R) \to \mathrm{H}^n(A_i)$ and $\beta_i^n : \mathrm{H}^n(A_i) \to \mathrm{H}^n(C)$ are compatible with the Yoneda products and the Lie brackets.

Proof. It is easily verified that the maps α^n and β^n of the above Proposition commute with the differential δ^n of Proposition 2.1.2 (b), giving rise to a short exact sequence of complexes

$$0 \longrightarrow \mathrm{Hom}_{E^e}(R^{\otimes\bullet}, M) \xrightarrow{\alpha^\bullet} \bigoplus_{i=1}^{2} \mathrm{Hom}_{E_i^e}(A_i^{\otimes\bullet}, e_i M e_i) \xrightarrow{\beta^\bullet} \mathrm{Hom}_{E_C^e}(C^{\otimes\bullet}, eMe) \longrightarrow 0$$

whose homology groups are the desired Hochschild cohomology groups. This shows the existence of the long exact sequence. In order to prove the second part of the statement, we need to show that $\alpha^\bullet = (\alpha_1^\bullet, \alpha_2^\bullet)^t$ is compatible with the cup-product and the Lie bracket if $M = R$. For the cup-product, this follows from the fact that if $x \in A_i^{\otimes_{E_i}(n+m)}$, then $x = e_i x e_i$ and

$$(\alpha_i^n(f)) \cup (\alpha_i^m(g))(x) = (f \cup g)(x) = \alpha_i^{n+m}(f \cup g)(x),$$

for $f \in \mathrm{Hom}_{E^e}(R^{\otimes_E n}, R)$ and $g \in \mathrm{Hom}_{E^e}(R^{\otimes_E m}, R)$. A similar argument applies to $\beta_i^\bullet$. For the Lie bracket, let $f \in \mathrm{Hom}_{E^e}(R^{\otimes_E n}, R)$, $g \in \mathrm{Hom}_{E^e}(R^{\otimes_E m}, M)$ and $x_1 \otimes \cdots \otimes x_{n+m-1} \in A_i^{\otimes n+m-1}$. Then

$$\begin{aligned}
&\alpha_i^{n+m-1}(f \circ_j g)(x_1 \otimes \cdots \otimes x_{n+m-1}) \\
&\qquad = (f \circ_j g)(e_i x_1 \otimes \cdots \otimes x_{n+m-1} e_i) \\
&\qquad = f(e_i x_1 \otimes \cdots \otimes g(x_j \otimes \cdots \otimes x_{j+m-1}) \otimes \cdots \otimes x_{n+m-1} e_i) \\
&\qquad = f(e_i x_1 \otimes \cdots \otimes g(e_i x_j \otimes \cdots \otimes x_{j+m-1} e_i) \otimes \cdots \otimes x_{n+m-1} e_i) \\
&\qquad = (\alpha_i^n(f)) \circ_j (\alpha_i^m(g))(x_1 \otimes \cdots \otimes x_{n+m-1}).
\end{aligned}$$

The result follows by linearity. Similarly, $\beta_i^\bullet$ is compatible with the Lie bracket. □

2.4. The case where C is a core algebra. This section is devoted to the study of a particular class of oriented bound quiver algebras.

More precisely, let $R = kQ_R/I_R$ where Q_R is a finite quiver and I_R is an admissible ideal in kQ_R. Moreover, the idempotents e, e'_1 and e'_2 are sums of primitive idempotents ε_x corresponding to some vertices x in Q_R, and Q_C, Q_{A_1} and Q_{A_2} are the full and convex subquivers of Q_R generated by these vertices. Finally, we have $I_C = kQ_C \cap I_R$, $I_1 = kQ_{A_1} \cap I_R$ and $I_2 = kQ_{A_2} \cap I_R$. Thus, $C = kQ_C/I_C$, $A_1 = kQ_{A_1}/I_1$ and $A_2 = kQ_{A_2}/I_2$ are also bound quiver algebras. For instance, the oriented algebra of Examples 2.5.2 (a) below is like so.

We show that if R is an oriented algebra as above and C satisfies some specific conditions, then the long exact sequence of Theorem 2.3.2 results in concrete formulæ to compute the Hochschild cohomology groups of R. The conditions we impose on $C = kQ_C/I_C$ are the following:

(1) Q_C contains no oriented cycles, and
(2) $\mathrm{H}^n(C) = 0$ for $n \geq 1$.

In the sequel, an algebra satisfying the above two conditions is called a **core algebra**. There exists many important classes of algebras which are core algebras, mostly related to trees.

Examples 2.4.1.

(a) Let $A = kQ/I$, where I is a two-sided **monomial** ideal and Q is connected (for instance $A = kQ$, a hereditary connected algebra). Then A is core if and only if Q is a tree, by [**16**, (1.6),(2.2)]

(b) We recall from [**16**] that an algebra A is called **piecewise hereditary of type** H if there exists a hereditary algebra H such that the bounded derived categories of finitely generated A-modules and H-modules are equivalent as triangulated categories. In this case, it follows from [**16**, (4.2)] that $\mathrm{H}^n(A)$ is isomorphic to $\mathrm{H}^n(H)$ for $n \geq 0$. Moreover, since, by [**15**, (IV.1.10)] the quiver of any piecewise hereditary algebra has no oriented cycles, then it follows from Example (a) that A is a core algebra if and only if the quiver of H is a tree.

(c) Following [**11**], for an algebra $A = kQ/I$, we say that $\mathrm{H}^n(A)$ **strongly vanishes** (see also [**26**, 2.2, 4.1] if $\mathrm{H}^n(B) = 0$ for every full convex subcategory B of A. Also, we say that A has **convex projectives** (or **convex injectives**) if for every vertex x of Q, the set of vertices $\{y \mid \varepsilon_x A \varepsilon_y \neq 0\}$ (or $\{y \mid \varepsilon_y A \varepsilon_x \neq 0\}$, respectively) is convex in Q. With these definitions, the following holds true by [**11**] : A is a core algebra provided
   (i) A is schurian and $\mathrm{H}^1(A)$ strongly vanishes, or
   (ii) Q has no oriented cycles and A is a tame algebra with convex projectives and convex injectives such that $\mathrm{H}^1(A)$ and $\mathrm{H}^2(A)$ strongly vanish.

We continue with the following lemma concerning the center of an algebra, whose proof is left as an exercise. Recall that the 0th Hochschild cohomology group $\mathrm{H}^0(A)$ of an algebra A is isomorphic to its center $Z(A)$.

Lemma 2.4.2. *Let $A = kQ/I$ be a connected algebra, ε_i be the idempotent corresponding to the vertex i, and ω be an element of the center $Z(A)$ of A. Then, $\omega = \sum_{i\in Q_0} \omega_i + \lambda \cdot 1_A$, where $\lambda \in k$ and $\omega_i \in \varepsilon_i \mathrm{rad} A \varepsilon_i$ for each i.*

Remark 2.4.3. It follows from the previous lemma that if $C = kQ_C/I_C$ is an algebra such that Q_C contains no oriented cycles, then $\dim_k \mathrm{H}^0(C)$ equals the number of connected components of Q_C.

We can now prove the main result of this section.

Proposition 2.4.4. *Let $R = kQ_R/I_R$ be a bound quiver algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = kQ_C/I_C$, $A_1 = kQ_{A_1}/I_1$ and $A_2 = kQ_{A_2}/I_2$. If $C = kQ_C/I_C$ is a core algebra, then:*
   (a) $\dim_k \mathrm{H}^0(R) = \dim_k \mathrm{H}^0(A_1) + \dim_k \mathrm{H}^0(A_2) - 1$;
   (b) $\dim_k \mathrm{H}^1(R) = \dim_k \mathrm{H}^1(A_1) + \dim_k \mathrm{H}^1(A_2) + \dim_k \mathrm{H}^0(C) - 1$;
   (c) $\mathrm{H}^i(R) \simeq \mathrm{H}^i(A_1) \oplus \mathrm{H}^i(A_2)$ *for* $i \geq 2$.

Proof. (a). It is sufficient to show that $\dim_k Z(R) = \dim_k Z(A_1) + \dim_k Z(A_2) - 1$. For $j = 1, 2$, let $\mathbb{B}_j$ be a k-basis of $Z(A_j)$ containing e_j, the unit element of A_j. We claim that $\mathbb{B} = (1_R \cup \mathbb{B}_1 \cup \mathbb{B}_2) \setminus (\{e_1\} \cup \{e_2\})$ is a k-basis of $Z(R)$. Since this set is linearly independent by Lemma 2.4.2, it remains to show that it generates $Z(R)$. Let ω be an element of $Z(R)$. By Lemma 2.4.2, we have $\omega = \sum_{i\in(Q_R)_0} \omega_i + \lambda \cdot 1_R$, where $\lambda \in k$ and $\omega_i \in \varepsilon_i \mathrm{rad} R \varepsilon_i$ for each $i \in (Q_R)_0$. Moreover, since Q_C has no oriented cycles, we have $\omega = \sum_{i\in(Q_R)_0 \setminus (Q_C)_0} \omega_i + \lambda \cdot 1_R$. Therefore, we can write without ambiguity $\omega = \rho_1 + \rho_2 + \lambda \cdot 1_R$, where $\rho_j = \sum_{i\in(Q_{A'_j})_0} \omega_i$, $j = 1, 2$. It then remains to show that $\rho_j \in Z(A_j)$ for $j = 1, 2$ since, by construction, $\rho_j \in Z(A_j)$ if and only if ρ_j belongs to the k-vector space generated by $\mathbb{B}_j \setminus \{e_j\}$. Let η_1 be a path in Q_{A_1}. We have :

$$\begin{aligned} \rho_1\eta_1 + \lambda\eta_1 &= \rho_1\eta_1 + 0 + \lambda\eta_1 = \rho_1\eta_1 + \rho_2\eta_1 + \lambda \cdot 1_R\eta_1 = \omega\eta_1 \\ &= \eta_1\omega = \eta_1\rho_1 + \eta_1\rho_2 + \lambda\eta_1 \cdot 1_R = \eta_1\rho_1 + 0 + \lambda\eta_1 \\ &= \eta_1\rho_1 + \lambda\eta_1 \end{aligned}$$

whence $\rho_1\eta_1 = \eta_1\rho_1$. Therefore, $\rho_1 \in Z(A_1)$. Since, similarly, $\rho_2 \in Z(A_2)$, ω belongs to the k-vector space generated by $\mathbb{B}$ and $\mathbb{B}$ is a k-basis of $Z(R)$. In particular, $\dim_k Z(R) = \dim_k Z(A_1) + \dim_k Z(A_2) - 1$, and this proves (a).

(b). Since $\mathrm{H}^i(C) = 0$ for each $i \geq 1$, it follows from Theorem 2.3.2 that we have the following exact sequence :

$$0 \longrightarrow \mathrm{H}^0(R) \longrightarrow \mathrm{H}^0(A_1) \oplus \mathrm{H}^0(A_2) \longrightarrow \mathrm{H}^0(C) \longrightarrow \mathrm{H}^1(R) \longrightarrow \mathrm{H}^1(A_1) \oplus \mathrm{H}^1(A_2) \longrightarrow 0$$

whence,

$$\begin{aligned} \dim_k\mathrm{H}^1(R) = \quad & \dim_k\mathrm{H}^1(A_1) + \dim_k\mathrm{H}^1(A_2) + \dim_k\mathrm{H}^0(C) \\ & + \dim_k\mathrm{H}^0(R) - \dim_k\mathrm{H}^0(A_1) - \dim_k\mathrm{H}^0(A_2) \end{aligned}$$

But $\dim_k\mathrm{H}^0(R) - \dim_k\mathrm{H}^0(A_1) - \dim_k\mathrm{H}^0(A_2) = -1$ by (a).

(c). This follows directly from Theorem 2.3.2. □

2.5. Examples and comparison. We now compute some examples and compare the sequence of Theorem 2.3.2 with other ones existing in the literature. First, let

$$R = \begin{pmatrix} A & 0 \\ {}_BM_A & B \end{pmatrix}$$

where A and B are arbitrary finitely generated k-algebras and M is a finitely generated B-A-bimodule. Then R is a k-algebra under the usual matrix operations.

In the eighties, D. Happel proved in [**16**] the existence of a long exact sequence relating the Hochschild cohomology groups of R, A and B in the particular case where $B = k$. Over the years, several variations and generalizations of this sequence have been obtained. The most recent long exact sequences now apply for an arbitrary algebra B (see [**9, 23, 13, 14, 2**]).

Theorem 2.5.1. *Let R be as above, e_A and e_B are the unit elements of A and B and let X be a R-R-bimodule. Then, there exists a long exact sequence*

$$0 \to \mathrm{H}^0(R, X) \to \mathrm{H}^0(A, e_AXe_A) \oplus \mathrm{H}^0(B, e_BXe_B) \to \mathrm{Ext}^0_{B\otimes_k A^{op}}(M, e_BXe_A) \to$$

$$\to \mathrm{H}^1(R, X) \to \mathrm{H}^1(A, e_AXe_A) \oplus \mathrm{H}^1(B, e_BXe_B) \to \mathrm{Ext}^1_{B\otimes_k A^{op}}(M, e_BXe_A) \to \cdots$$

The following examples show the difference between the exact sequence of Theorem 2.3.2 and the above sequence.

Examples 2.5.2.

(a) Let R be the bound quiver algebra given by the quiver



bounded by the ideal generated by all paths of length two. Let $e = \varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4$, $e'_1 = \varepsilon_5$, and $e'_2 = \varepsilon_6$ be the three idempotents orienting R. By [**8**, (p. 96)],

$$\mathrm{H}^i(A_1) \simeq \mathrm{H}^i(A_2) \simeq \begin{cases} k, & \text{if } i = 0; \\ k^2, & \text{if } i = 1; \\ 0, & \text{otherwise;} \end{cases} \quad \text{and} \quad \mathrm{H}^i(R) \simeq \begin{cases} k, & \text{if } i = 0; \\ k^4, & \text{if } i = 1; \\ 0, & \text{otherwise.} \end{cases}$$

Moreover, since the quiver of C is a tree, $\mathrm{H}^0(C) = k$ and $\mathrm{H}^i(C) = 0$ for $i \geq 1$. By Theorem 2.3.2 and Proposition 2.4.4, there exists an exact sequence :

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{H}^0(R) \arrow[r] \arrow[d, equal] & \mathrm{H}^0(A_1)\oplus\mathrm{H}^0(A_2) \arrow[r] \arrow[d, equal] & \mathrm{H}^0(C) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(R) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(A_1)\oplus\mathrm{H}^1(A_2) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(C) \arrow[r] \arrow[d, equal] & \cdots \\
0 \arrow[r] & \boxed{k} \arrow[r] & k\oplus k \arrow[r] & k \arrow[r] & \boxed{k^4} \arrow[r] & k^2\oplus k^2 \arrow[r] & 0 \arrow[r] & \cdots
\end{tikzcd}$$

On the other hand, if we respectively denote by S_x and P_x the simple module and the projective module associated to the vertex x in the quiver, one can see R as the one-point extension

$$A_1[M] = \begin{pmatrix} A_1 & 0 \\ M & k \end{pmatrix},$$

where $M = \mathrm{rad} P_6 = S_2 \oplus S_3 \oplus S_4$ is a semisimple module without self-extensions. In this case, the long exact sequence of Theorem 2.5.1 is given by :

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{H}^0(R) \arrow[r] \arrow[d, equal] & \mathrm{H}^0(A_1)\oplus k \arrow[r] \arrow[d, equal] & \mathrm{Hom}_{A_1}(M,M) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(R) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(A_1) \arrow[r] \arrow[d, equal] & \mathrm{Ext}^1_{A_1}(M,M) \arrow[r] \arrow[d, equal] & \cdots \\
0 \arrow[r] & k \arrow[r] & k\oplus k \arrow[r] & k^3 \arrow[r] & k^4 \arrow[r] & k^2 \arrow[r] & 0 \arrow[r] & \cdots
\end{tikzcd}$$

The two long exact sequences are clearly different.

(b) Let R be the algebra given by the quiver



with the relations $\alpha^2 = \alpha\beta = \beta\gamma = \delta\varepsilon = \varepsilon^2 = 0$. Let $e = \varepsilon_3$, $e'_1 = \varepsilon_1 + \varepsilon_2$ and $e'_2 = \varepsilon_4$. Then the conditions (1.1) and (1.4) of Definition 1.2.2 are satisfied, and so R is oriented. A quick application of [**8**, (p. 96)] and Theorem 2.5.1 gives:

$$\mathrm{H}^i(A_1) \simeq \mathrm{H}^i(A_2) \simeq \begin{cases} k^2, & \text{if } i = 0; \\ k, & \text{otherwise .} \end{cases}$$

Moreover, $\mathrm{H}^0(C) = k$ and $\mathrm{H}^i(C) = 0$ for $i \geq 1$. By Theorem 2.3.2 and Proposition 2.4.4, there exists an exact sequence

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{H}^0(R) \arrow[r] \arrow[d, equal] & \mathrm{H}^0(A_1)\oplus\mathrm{H}^0(A_2) \arrow[r] \arrow[d, equal] & \mathrm{H}^0(C) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(R) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(A_1)\oplus\mathrm{H}^1(A_2) \arrow[r] \arrow[d, equal] & \mathrm{H}^1(C) \arrow[r] \arrow[d, equal] & \cdots \\
0 \arrow[r] & \boxed{k^3} \arrow[r] & k^2\oplus k^2 \arrow[r] & k \arrow[r] & \boxed{k^2} \arrow[r] & k\oplus k \arrow[r] & 0 \arrow[r] & \cdots
\end{tikzcd}$$

from which we obtain that $\mathrm{H}^1(R) \simeq k^2$ and $\mathrm{H}^i(R) \simeq \mathrm{H}^i(A_1) \oplus \mathrm{H}^i(A_2) \simeq k \oplus k$ for $i \geq 2$. On the other hand, one can see R as the following matrix algebra

$$R = \begin{pmatrix} A_1 & 0 \\ M & A'_2 \end{pmatrix},$$

where A_2' is the full subcategory generated by the vertex 4 and M is the A_2'-A_1-bimodule given by the arrow β. In order to apply the sequence of Theorem 2.5.1, one needs to compute $\mathrm{Ext}^i_{A_2'\otimes {A_1}^{op}}(M,M)$ for $i \geq 0$. A direct computation leads to the fact that $A_2' \otimes {A_1}^{op}$ is given by the quiver

$$\begin{tikzcd}
4\otimes 1 \arrow[loop above, "4\otimes\varepsilon"] \arrow[loop below, "\alpha\otimes 1"'] \arrow[r, "4\otimes\delta"] & 4\otimes 2 \arrow[loop below, "\alpha\otimes 2"'] \arrow[r, "4\otimes\gamma"] & 4\otimes 3 \arrow[loop below, "\alpha\otimes 3"']
\end{tikzcd}$$

bound by the relations induced by those of R, that is $(\alpha\otimes 1)^2 = (\alpha\otimes 2)^2 = (\alpha\otimes 3)^2 = (4\otimes\varepsilon)^2 = (4\otimes\varepsilon)(4\otimes\delta) = 0$, $(\alpha\otimes 1)(4\otimes\delta) = (4\otimes\delta)(\alpha\otimes 2)$ and $(\alpha\otimes 2)(4\otimes\gamma) = (4\otimes\gamma)(\alpha\otimes 3)$. Moreover, the $A_2'\otimes {A_1}^{op}$-module M is the simple module $S_{4\otimes 3}$. However, since $(\alpha\otimes 3)^2 = 0$, it follows from [**19**, (1.4)] that $\mathrm{Ext}^i_{A_2'\otimes {A_1}^{op}}(M,M) \neq 0$ for $i \geq 1$. Hence, the long exact sequence of Theorem 2.5.1 cannot efficiently be used to compute the Hochschild cohomology groups of R. At least, it cannot be used to show that $\mathrm{H}^i(R) \simeq \mathrm{H}^i(A_1) \oplus \mathrm{H}^i(A_2) \simeq k \oplus k$ for $i \geq 2$.

3. Cyclic (co)homology

The strong relation between cyclic (co)homology and Hochschild (co)homology theories brings up the natural question whether oriented algebras give rise to long exact sequences of cyclic (co)homology groups. In what follows, we show that the answer is positive, and moreover that these long exact sequences are compatible with Connes' long exact sequences.

The first part of this section is devoted to basic definitions and tools concerning cyclic (co)homology, including the Connes' long exact sequences. Then we consider the oriented algebras again and prove our main results.

We refer to [**20**] and [**3**] for more details on cyclic (co)homology theory.

3.1. Definitions and notations. We first recall the following general constructions. A **bicomplex** $C_{\bullet\bullet}$ is a collection of modules C_{pq}, with $p, q \in \mathbb{Z}$, together with a horizontal differential $d^h : C_{p,q} \to C_{p-1,q}$ and a vertical differential $d^v : C_{p,q} \to C_{p,q-1}$ satisfying $d^v d^v = d^h d^h = d^v d^h + d^h d^v = 0$. Now, given a bicomplex $C_{\bullet\bullet}$ in the first quadrant, that is $C_{p,q} = 0$ provided $p < 0$ or $q < 0$, define, for $n \in \mathbb{Z}$, the module $(\mathrm{Tot} C_{\bullet\bullet})_n = \bigoplus_{p+q=n} C_{p,q}$, and, further, the complex

$$\cdots \longrightarrow (\mathrm{Tot}C_{\bullet\bullet})_{n+1} \xrightarrow{d} (\mathrm{Tot}C_{\bullet\bullet})_n \xrightarrow{d} (\mathrm{Tot}C_{\bullet\bullet})_{n-1} \longrightarrow \cdots$$

where $d = d^h + d^v$. This complex is called the **total complex** of the bicomplex $C_{\bullet\bullet}$. Then it is a routine verification that a short exact sequence of bicomplexes in the first quadrant yields a short exact sequence of their associated total bicomplexes.

Now, let R be an algebra. For each $n \geq 2$, consider the morphism of R-bimodules $b : R^{\otimes n} \to R^{\otimes n-1}$ given by

$$\begin{aligned} (3.1) \qquad b(r_0 \otimes \cdots \otimes r_{n-1}) \quad := \quad & \sum_{i=0}^{n-2} (-1)^i (r_0 \otimes r_1 \otimes \cdots \otimes r_i r_{i+1} \otimes \cdots \otimes r_{n-1}) \\ & + (-1)^{n-1} (r_{n-1} r_0 \otimes r_1 \otimes \cdots \otimes r_{n-2}) \end{aligned}$$

It is easily verified that $b \circ b = 0$, so we get a complex $(R^{\otimes \bullet}, b)$.

On the other hand, for $n \geq 1$, the cyclic group $\mathbb{Z}_{n+1}$ acts on $R^{\otimes n+1}$ by letting its generator t acts by $t(r_0 \otimes \cdots \otimes r_n) := (-1)^n (r_n \otimes r_0 \cdots \otimes r_{n-1})$. For $n = 0$, let t be the identity map. So t defines an endomorphism on $R^{\otimes n+1}$. Let $N = 1 + t + \cdots + t^n$. Then the following diagram

(3.2)
$$\begin{tikzcd}
\vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \\
R^{\otimes 3} \arrow[d, "b"'] & R^{\otimes 3} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & R^{\otimes 3} \arrow[l, "N"'] \arrow[d, "b"'] & R^{\otimes 3} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & \cdots \arrow[l, "N"'] \\
R^{\otimes 2} \arrow[d, "b"'] & R^{\otimes 2} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & R^{\otimes 2} \arrow[l, "N"'] \arrow[d, "b"'] & R^{\otimes 2} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & \cdots \arrow[l, "N"'] \\
R & R \arrow[l, "1-t"'] & R \arrow[l, "N"'] & R \arrow[l, "1-t"'] & \cdots \arrow[l, "N"']
\end{tikzcd}$$

where b and b' are as in (3.1) and (2.1), is a bicomplex in the first quadrant. This diagram is called the **cyclic bicomplex** associated with R and is denoted $CC(R)$.

By definition, the nth **cyclic homology group** of R is given by

$$\mathrm{HC}_n(R) := \mathrm{H}_n(\mathrm{Tot}CC(R)),$$

that is the nth homology group of the total complex associated with $CC(R)$.

In order to define the cyclic cohomology groups, we consider the standard duality $D = \mathrm{Hom}_k(-, k)$. Applying D on the bicomplex $CC(R)$ above yields the dual bicomplex $D(CC(R))$. By definition, the nth **cyclic cohomology group** $\mathrm{HC}^n(R)$ of R is the nth homology group of the cochain complex $\mathrm{Tot}D(CC(R))$:

$$\mathrm{HC}^n(R) := \mathrm{H}^n(\mathrm{Tot}D(CC(R))).$$

3.2. Connes' exact sequences. Let $CC(R)^{\{2\}}$ be the bicomplex consisting of the first two columns of $CC(R)$. By Killing's lemma (see [**20**, p. 55] for instance) the homology groups of $\mathrm{Tot}CC(R)^{\{2\}}$ are isomorphic to the homology groups of the first column, which are the Hochschild homology group $\mathrm{H}_n(R)$ because $1_R \otimes b'$ becomes b under the isomorphism $R \otimes_{R^e} R^{\otimes n+2} \simeq R \otimes R^{\otimes n}$ given by $r_0 \otimes r_1 \otimes \cdots \otimes r_{n+2} \mapsto r_{n+2} r_0 r_1 \otimes r_2 \otimes \cdots \otimes r_{n+1}$. Now, let $(CC(R)[2,0])_{pq} = CC(R)_{p-2,q}$. Thus, the short exact sequence of bicomplexes

$$0 \longrightarrow CC(R)^{\{2\}} \longrightarrow CC(R) \longrightarrow CC(R)[2,0] \longrightarrow 0$$

yields a short exact sequence of total complexes, and so a long exact sequence

$$\cdots \longrightarrow \mathrm{H}_n(R) \longrightarrow \mathrm{HC}_n(R) \longrightarrow \mathrm{HC}_{n-2}(R) \longrightarrow \mathrm{H}_{n-1}(R) \longrightarrow \cdots$$

relating the Hochschild homology groups and cyclic homology groups of R, called the **homological Connes' exact sequence**.

Dually, let $D(CC(R))^{\{2\}}$ denote the bicomplex consisting of the first two columns of $D(CC(R))$. Since D is an exact functor, Killing's lemma applies and the homology groups of $\mathrm{Tot}D(CC(R))^{\{2\}}$ are isomorphic to the homology groups of the first column, which are the Hochschild cohomology groups $\mathrm{H}^n(R, D(R))$ because $\mathrm{Hom}_{R^e}(b', D(R))$ becomes $D(b)$ under the isomorphism $\phi : \mathrm{Hom}_{R^e}(R^{\otimes n+1}, D(R)) \to \mathrm{Hom}_k(R^{\otimes n}, k)$ sending any $f \in \mathrm{Hom}_{R^e}(R^{\otimes n+1}, D(R))$ to the map $\phi(f)$ defined by

$\phi(f)(r_0 \otimes \cdots \otimes r_{n-1}) = f(1 \otimes r_1 \otimes \cdots \otimes r_{n-1} \otimes 1)(r_0)$ for $r_0 \otimes \cdots \otimes r_{n-1} \in R^{\otimes n}$. Therefore, the short exact sequence of bicomplexes

$$\begin{tikzcd} 0 \arrow[r] & D(CC(R))[2,0] \arrow[r] & D(CC(R)) \arrow[r] & D(CC(R))^{\{2\}} \arrow[r] & 0 \end{tikzcd}$$

(where $D(CC(R)[2,0])_{pq} = D(CC(R))_{p-2,q}$) yields a short exact sequence of total complexes giving rise to the **cohomological Connes' exact sequence**

$$\cdots \longrightarrow \mathrm{H}^n(R, D(R)) \longrightarrow \mathrm{HC}^{n-1}(R) \longrightarrow \mathrm{HC}^{n+1}(R) \longrightarrow \mathrm{H}^{n+1}(R, D(R)) \longrightarrow \cdots$$

3.3. Cyclic (co)homology for oriented algebras. Let R be an algebra, oriented by three idempotents e, e_1', e_2'. We also adopt the notations and terminology of Section 1.2. In what follows, we give an analogues to Theorems 2.2.3 and 2.3.2 for the cyclic (co)homology groups.

The strategy is the same as for Hochschild (co)homology groups. Thus, in view of Remark 2.2.2, it will be necessary to deal with tensor products over E rather than over k. This means that it would be nice to construct a short exact sequence relating the bicomplexes $CC_E(C)$, $CC_E(A_1)$, $CC_E(A_2)$ and $CC_E(R)$ (where these are obtained from the bicomplexes (3.2) by replacing the tensor products over k by tensor products over E), giving rise to a long exact sequence of homology groups. However, although such a sequence exists, this is unfortunately not sufficient because there is no guarantee, for instance, that the total complexes of $CC(R)$ and $CC_E(R)$ have the same homology groups. This problem can be fixed by what is sometimes called the relative Hochschild homology.

Let S be a subring of R (in our case we will take $S = E$). In particular, R is a S-bimodule. Consider the S-subbimodule J^n of $R^{\otimes sn}$ generated by all elements of the form $(sr_1 \otimes r_2 \otimes \cdots \otimes r_n) - (r_1 \otimes r_2 \otimes \cdots \otimes r_n s)$ for any $s \in S$ and $r_i \in R$. Given $x, x' \in R^{\otimes sn}$, we write

$$x \sim x' \text{ provided } x - x' \in J^n, \text{ and define } \widetilde{R^{\otimes sn}} := R^{\otimes sn}/J^n.$$

It is straightforward to check that the differential b of (3.1) is compatible with the relation $\sim$, in the sense that if $x \sim x'$ then $b(x) \sim b(x')$. So there is a well-defined complex $(\widetilde{R^{\otimes sn}}, b)$. It's nth homology group is denoted $\widetilde{\mathrm{H}_n^S}(R)$.

Recall that an algebra S is called **separable** over k if the S-bimodule map $\mu : S \otimes S^{op} \to S$ splits. This is equivalent to the existence of an idempotent $\varepsilon = \sum u_i \otimes v_i \in S \otimes S^{op}$ such that $\sum u_i v_i = 1$ and $(s \otimes 1_S)\varepsilon = (1_S \otimes s)\varepsilon$ for any $s \in S$.

Then we have the following theorem from [**20**, (p.21)].

THEOREM 3.3.1. *Let S be separable over k. Then the canonical projection $\pi : (R^{\otimes n}, b) \to (\widetilde{R^{\otimes sn}}, b)$ is a quasi-isomorphism. In particular,* $\mathrm{H}_n(R) \simeq \widetilde{\mathrm{H}_n^S}(R)$ *for any $n \geq 0$.*

Moreover, it is straightforward to check that if, in the diagram (3.2), we replace the tensor products over k by tensors products over S, to obtain a complex $CC_S(R)$, the induced applications b, $-b'$, t and N are compatible with the relation $\sim$, so that we obtain a bicomplex $\widetilde{CC_S}(R)$, refered in the sequel as the **relative bicomplex**, given by

(3.3)
$$\begin{tikzcd}
\vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \\
\widetilde{R^{\otimes_S 3}} \arrow[d, "b"'] & \widetilde{R^{\otimes_S 3}} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & \widetilde{R^{\otimes_S 3}} \arrow[l, "N"'] \arrow[d, "b"'] & \widetilde{R^{\otimes_S 3}} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & \cdots \arrow[l, "N"'] \\
\widetilde{R^{\otimes_S 2}} \arrow[d, "b"'] & \widetilde{R^{\otimes_S 2}} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & \widetilde{R^{\otimes_S 2}} \arrow[l, "N"'] \arrow[d, "b"'] & \widetilde{R^{\otimes_S 2}} \arrow[l, "1-t"'] \arrow[d, "-b'"'] & \cdots \arrow[l, "N"'] \\
\widetilde{R} & \widetilde{R} \arrow[l, "1-t"'] & \widetilde{R} \arrow[l, "N"'] & \widetilde{R} \arrow[l, "1-t"'] & \cdots \arrow[l, "N"']
\end{tikzcd}$$

and the map π of Theorem 3.3.1 extends to a map of bicomplexes $\pi : CC(R) \to \widetilde{CC_S}(R)$. Moreover, with these notations, Theorem 3.3.1 assures that the restriction of π to the columns of type b is a quasi-isomorphism. On the other hand, since the columns of type b' are exact, the projection π trivially induces a quasi-isomorphism on the columns of type b'. Then, it follows (see [**20**, Proposition 1.0.12] for instance) that the map induced by π on the total complexes is a quasi-isomorphism. We have just proved the following proposition.

Proposition 3.3.2. *Let R be an algebra and S be a separable subalgebra of R. Then* $\mathrm{HC}_n(R) \simeq \mathrm{H}_n(\mathrm{Tot}\widetilde{CC_S}(R))$ *for each* $n \geq 0$.

For an oriented algebra R, the subalgebra $S = E$ of R is separable over k (take $\varepsilon = e'_1 \otimes e'_1 + e \otimes e + e'_2 \otimes e'_2$), and so are the subalgebras E_C of C, E_1 of A_1 and E_2 of A_2. As a consequence, to obtain a long exact sequence of cyclic homology groups, it suffices to construct a short exact sequence of relative bicomplexes of R, C, A_1 and A_2. The existence of such a sequence will follow from the next proposition.

Proposition 3.3.3. *Let R be an algebra, oriented by three idempotents e, e'_1, e'_2. For any $n \geq 1$, the sequence*

$$0 \longrightarrow C^{\otimes_{E_C} n} \xrightarrow{f^n} A_1^{\otimes_{E_1} n} \bigoplus A_2^{\otimes_{E_2} n} \xrightarrow{g^n} R^{\otimes_E n} \longrightarrow 0$$

is exact in $\mathrm{mod}E^e$, *where* $f^n = (f_1^n, f_2^n)^t$, *with* $f_i^n(er_1 \otimes \cdots \otimes r_n e) = e_i e r_1 \otimes \cdots \otimes r_n e e_i$, *and* $g^n = (g_1^n, g_2^n)$, *with* $g_i^n(e_i r_1 \otimes \cdots \otimes r_n e_i) = (-1)^i (e_i r_1 \otimes \cdots \otimes r_n e_i)$, *for* $i = 1, 2$.

Proof. Since $e_i e = e = e e_i$, the map f^n is trivially injective. Also, $\mathrm{Im} g^n \subseteq \mathrm{Ker} f^n$. Conversely, assume $x_1 = \sum_{i=1}^k e_1 r_{1i} \otimes \cdots \otimes r_{ni} e_1 \in A_1^{\otimes_E n}$ and $x_2 = \sum_{i=1}^l e_2 r'_{1i} \otimes \cdots \otimes r'_{ni} e_2 \in A_2^{\otimes_E n}$ are such that $g^n((x_1, x_2)) = 0$. By definition of g^n, we get $x_1 - x_2 = 0$, and so $x_1 = x_2$. Let x be this common value. Multiplying x by e_1 on both sides gives $x = x_1 = e_1 x_1 e_1 = e_1 x_2 e_1 = e_1 x_2 e_1 = \sum_{i=1}^l e r'_{1i} \otimes \cdots \otimes r'_{ni} e$ since $e_1 e_2 = e = e_2 e_1$. So $x \in C^{\otimes_E n}$ and $f^n(x) = (x_1, x_2)$, showing that $\mathrm{Ker} g^n \subseteq \mathrm{Im} f^n$. It remains to show that g^n is surjective. Let $r \in R^{\otimes_E n}$. Since $1_R = e'_1 + e + e'_2$, and $e'_1 r e'_2 = 0 = e'_2 r e'_1$ by Remark 1.3.1 (c), we have

$$\begin{aligned} r &= e'_1 r e'_1 + e'_1 r e + e'_1 r e'_2 + e r e'_1 + e r e + e r e'_2 + e'_2 r e'_1 + e'_2 r e + e'_2 r e'_2 \\ &= e'_1 r e'_1 + e'_1 r e + e r e'_1 + e r e + e r e'_2 + e'_2 r e + e'_2 r e'_2 \end{aligned}$$

Moreover, since $e_1e'_1 = e'_1 = e'_1e_1$, we have $e_1e = e = ee_1$. Similarly, $e_2e = e = ee_2$. We get $r_1 := e'_1re'_1 + e'_1re + ere'_1 + ere \in e_1R^{\otimes_E n}e_1 = A_1^{\otimes_E n}$ and $r_2 := ere'_2 + e'_2re + e'_2re'_2 \in e_2R^{\otimes_E n}e_2 = A_2^{\otimes_E n}$. This shows that $g^n(r_1, -r_2) = r$ and g^n is surjective. Similar arguments show that f^n and g^n are E^e-linear. □

Corollary 3.3.4. *Let R be an algebra, oriented by three idempotents e, e'_1, e'_2. For any $n \geq 1$, there exists a short exact sequence in* $\text{mod}E^e$ *of the form*

$$\begin{tikzcd}
0 \arrow[r] & \widetilde{C^{\otimes_{E_C} n}} \arrow[r, "f^n"] & \widetilde{A_1^{\otimes_{E_1} n}} \bigoplus \widetilde{A_2^{\otimes_{E_2} n}} \arrow[r, "g^n"] & \widetilde{R^{\otimes_E n}} \arrow[r] & 0
\end{tikzcd}$$

Proof. It suffices to check that the maps f^n and g^n in Proposition 3.3.3 are compatible with the relation $\sim$ and remain respectively injective and surjective, which is straightforward. □

We can now state and prove the main results of this section.

Theorem 3.3.5. *Let R be an algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_1 = e_2Re_2$ be as in Definition 1.2.2. There exists a long exact sequence of cyclic homology groups :*

$$\begin{tikzcd}
\cdots \arrow[r] & \mathrm{HC}_1(C) \arrow[r] & \mathrm{HC}_1(A_1) \oplus \mathrm{HC}_1(A_2) \arrow[r] & \mathrm{HC}_1(R) \arrow[r] & {} \\
{} \arrow[r] & \mathrm{HC}_0(C) \arrow[r] & \mathrm{HC}_0(A_1) \oplus \mathrm{HC}_0(A_2) \arrow[r] & \mathrm{HC}_0(R) \arrow[r] & 0
\end{tikzcd}$$

Proof. The maps f^n and g^n of Corollary 3.3.4 commute with b, b', t and N in (3.3), so they extend to a short exact sequence of relative bicomplexes

$$\begin{tikzcd}
0 \arrow[r] & \widetilde{CC_{E_C}}(C) \arrow[r, "f^\bullet"] & \widetilde{CC_{E_1}}(A_1) \oplus \widetilde{CC_{E_2}}(A_2) \arrow[r, "g^\bullet"] & \widetilde{CC_E}(R) \arrow[r] & 0
\end{tikzcd} \tag{3.4}$$

The result then follows from Proposition 3.3.2. □

In Theorem 2.2.3 and Theorem 3.3.5, we have obtained long exact sequences of Hochschild homology groups and cyclic homology groups. A deep inspection of the natural morphisms involved in these sequences, as well as in Connes' long exact sequence yields the following result, whose verification is left to the reader.

Theorem 3.3.6. *Let R be an algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_1 = e_2Re_2$ be as in Definition 1.2.2.*

There exists a commutative diagram

$$\begin{tikzcd}
 & \vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \\
\cdots \arrow[r] & \mathrm{H}_n(R) \arrow[r] \arrow[d] & \mathrm{HC}_n(R) \arrow[r] \arrow[d] & \mathrm{HC}_{n-2}(R) \arrow[r] \arrow[d] & \mathrm{H}_{n-1}(R) \arrow[r] \arrow[d] & \cdots \\
\cdots \arrow[r] & \mathrm{H}_{n-1}(C) \arrow[r] \arrow[d] & \mathrm{HC}_{n-1}(C) \arrow[r] \arrow[d] & \mathrm{HC}_{n-3}(C) \arrow[r] \arrow[d] & \mathrm{H}_{n-2}(C) \arrow[r] \arrow[d] & \cdots \\
\cdots \arrow[r] & \begin{array}{c}\mathrm{H}_{n-1}(A_1)\\ \oplus \\ \mathrm{H}_{n-1}(A_2)\end{array} \arrow[r] \arrow[d] & \begin{array}{c}\mathrm{HC}_{n-1}(A_1)\\ \oplus \\ \mathrm{HC}_{n-1}(A_2)\end{array} \arrow[r] \arrow[d] & \begin{array}{c}\mathrm{HC}_{n-3}(A_1)\\ \oplus \\ \mathrm{HC}_{n-3}(A_2)\end{array} \arrow[r] \arrow[d] & \begin{array}{c}\mathrm{H}_{n-2}(A_1)\\ \oplus \\ \mathrm{H}_{n-2}(A_2)\end{array} \arrow[r] \arrow[d] & \cdots \\
\cdots \arrow[r] & \mathrm{H}_{n-1}(R) \arrow[r] \arrow[d] & \mathrm{HC}_{n-1}(R) \arrow[r] \arrow[d] & \mathrm{HC}_{n-3}(R) \arrow[r] \arrow[d] & \mathrm{H}_{n-2}(R) \arrow[r] \arrow[d] & \cdots \\
 & \vdots & \vdots & \vdots & \vdots &
\end{tikzcd}$$

in which the rows and columns are exact.

We finish this section by showing that dual results hold for cyclic cohomology groups. Let R be an algebra and S be a separable subalgebra. We have seen (in our discussion before Proposition 3.3.2) that the canonical projection $\pi : CC(R) \to \widetilde{CC_S}(R)$ is a morphism of bicomplexes which is a quasi-isomorphism when restricted to the columns. Because $D = \mathrm{Hom}_k(-,k)$ is exact, the dual morphism $D(\pi) : D(\widetilde{CC_S}(R)) \to D(CC(R)$ has the same property. Thus the map induced by $D(\pi)$ on the total complexes is a quasi-isomorphism. We get the following.

Proposition 3.3.7. *Let R be an algebra and S be a separable subalgebra of R. Then* $\mathrm{HC}^n(R) \simeq \mathrm{H}^n(\mathrm{Tot}D(\widetilde{CC_S}(R)))$.

As a consequence, to obtain a long exact sequence on cyclic homology groups, it suffices to construct a short exact sequence of the duals of the relative bicomplexes (for $S = E$). But such a sequence exists by (3.4) and the fact that D is exact. We have shown the following theorem.

Theorem 3.3.8. *Let R be an algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_1 = e_2Re_2$ be as in Definition 1.2.2. There exists a long exact sequence of cyclic cohomology groups:*

$$0 \longrightarrow \mathrm{HC}^0(R) \longrightarrow \mathrm{HC}^0(A_1) \oplus \mathrm{HC}^0(A_2) \longrightarrow \mathrm{HC}^0(C) \longrightarrow$$

$$\longrightarrow \mathrm{HC}^1(R) \longrightarrow \mathrm{HC}^1(A_1) \oplus \mathrm{HC}^1(A_2) \longrightarrow \mathrm{HC}^1(C) \longrightarrow \cdots$$

Here again, the long exact sequences obtained in Theorem 2.3.2 and Theorem 3.3.8 embed with the cohomological Connes' exact sequence, so that we get the following result, whose verification is left to the reader.

Theorem 3.3.9. *Let R be an algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, let $C = eRe$, $A_1 = e_1Re_1$ and $A_1 = e_2Re_2$ be as in Definition 1.2.2.*

There exists a commutative diagram

$$\begin{tikzcd}
 & \vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \vdots \arrow[d] & \\
\cdots \arrow[r] & \mathrm{H}^n(R, D(R)) \arrow[r] \arrow[d] & \mathrm{HC}^{n-1}(R) \arrow[r] \arrow[d] & \mathrm{HC}^{n+1}(R) \arrow[r] \arrow[d] & \mathrm{H}^{n+1}(R, D(R)) \arrow[r] \arrow[d] & \cdots \\
\cdots \arrow[r] & \begin{matrix} \mathrm{H}^n(A_1, D(A_1)) \\ \oplus \\ \mathrm{H}^n(A_2, D(A_2)) \end{matrix} \arrow[r] \arrow[d] & \begin{matrix} \mathrm{HC}^{n-1}(A_1) \\ \oplus \\ \mathrm{HC}^{n-1}(A_2) \end{matrix} \arrow[r] \arrow[d] & \begin{matrix} \mathrm{HC}^{n+1}(A_1) \\ \oplus \\ \mathrm{HC}^{n+1}(A_2) \end{matrix} \arrow[r] \arrow[d] & \begin{matrix} \mathrm{H}^{n+1}(A_1, D(A_1)) \\ \oplus \\ \mathrm{H}^{n+1}(A_2, D(A_2)) \end{matrix} \arrow[r] \arrow[d] & \cdots \\
\cdots \arrow[r] & \mathrm{H}^n(C, D(C)) \arrow[r] \arrow[d] & \mathrm{HC}^{n-1}(C) \arrow[r] \arrow[d] & \mathrm{HC}^{n+1}(C) \arrow[r] \arrow[d] & \mathrm{H}^{n+1}(C, D(C)) \arrow[r] \arrow[d] & \cdots \\
\cdots \arrow[r] & \mathrm{H}^{n+1}(R, D(R)) \arrow[r] \arrow[d] & \mathrm{HC}^{n}(R) \arrow[r] \arrow[d] & \mathrm{HC}^{n+2}(R) \arrow[r] \arrow[d] & \mathrm{H}^{n+2}(R, D(R)) \arrow[r] \arrow[d] & \cdots \\
 & \vdots & \vdots & \vdots & \vdots &
\end{tikzcd}$$

in which the rows and columns are exact.

4. Simplicial (co)homology and fundamental group

In this section we consider algebras of the form $R = kQ_R/I_R$. More precisely we study the simplicial (co)homology groups of R (in case they admit semi-normed basis) as well as the fundamental group $\pi_1(Q_R, I_R)$. As before, we show the existence of Mayer-Vietoris long exact sequence relating the simplicial (co)homology groups. Then we prove that under some additional assumptions, it is possible to express the fundamental group of (Q_R, I_R) in terms of those of (Q_C, I_C), (Q_{A_1}, I_{A_1}) and (Q_{A_2}, I_{A_2}). We further apply this result to the computation of the first Hochschild cohomology group of some schurian oriented algebras.

We stress here that the validity of one of the conditions (1.2)-(1.4) is not necessary in the sequel, and so the results contained in this section apply for a slightly more general family of algebras, that is those satisfying (possibly only) condition (1.1), for instance the algebra of Examples 1.2.1 (b). Nevertheless, we will not make any distinction and we will keep going with the expression "oriented algebras". Also, all algebras considered in this section are bound quiver algebras, that is of the form kQ/I for some bound quiver (Q, I) (see Sections 1.1 and 2.4 for more details).

4.1. Simplicial (co)homology. Given an algebra $R = kQ/I$ and a pair (x, y) of vertices of Q, let ${}_y\mathbb{B}_x$ be a basis of the k-space $\varepsilon_x R \varepsilon_y$ and $\mathbb{B} = \bigcup_{(x,y)\in Q_0\times Q_0} {}_y\mathbb{B}_x$. We say that $\mathbb{B}$ is a **semi-normed basis** [**21**] if:

(a) $\varepsilon_x \in {}_x\mathbb{B}_x$ for every vertex $x \in Q_0$;
(b) $\alpha + I \in {}_y\mathbb{B}_x$ for every arrow $\alpha : x \to y$;
(c) For σ and σ' in $\mathbb{B}$, either $\sigma\sigma' = 0$ or there exists $0 \neq \lambda_{\sigma,\sigma'} \in k$, and $b(\sigma, \sigma') \in \mathbb{B}$ such that $\sigma\sigma' = \lambda_{\sigma,\sigma'} b(\sigma, \sigma')$.

Assume $R = kQ_R/I_R$ admits a semi-normed basis. Following [**22, 21**], one can associate to R the chain complex $(\mathrm{SC}_\bullet(R), d)$ as follows: $\mathrm{SC}_0(R)$ and $\mathrm{SC}_1(R)$ are the free abelian groups with basis Q_0 and $\mathbb{B}$, respectively. For $n \geq 2$, let

$\mathrm{SC}_n(R)$ be the free abelian group with basis the set of n-tuples $(\sigma_1, \sigma_2, \dots, \sigma_n)$ of $\mathbb{B}^n$ such that $\sigma_1\sigma_2\cdots\sigma_n \neq 0$, and $\sigma_i \neq e_j$ for all $i, j \in Q_0$. The differential $d_n : \mathrm{SC}_n(R) \to \mathrm{SC}_{n-1}(R)$ is defined on the basis elements by the rules $d_1(\sigma) = y - x$ for $\sigma \in {}_y\mathbb{B}_x$ and

$$\begin{aligned} d_n(\sigma_1, \sigma_2, \dots, \sigma_n) &= (\sigma_2, \dots, \sigma_n) \\ &+ \sum_{j=1}^{n-1} (-1)^j (\sigma_1, \dots, b(\sigma_j, \sigma_{j+1}), \dots, \sigma_n) \\ &+ (-1)^n (\sigma_1, \dots, \sigma_{n-1}) \end{aligned}$$

for $n \geq 1$ and $(\sigma_1, \sigma_2, \dots, \sigma_n) \in \mathrm{SC}_n(R)$.

The nth **simplicial homology group** of R (with respect to the basis $\mathbb{B}$) is the nth homology group of the chain complex $(\mathrm{SC}_\bullet(R), d_\bullet)$, and we denote it by $\mathrm{SH}_n(R)$. Given an abelian group G, the nth-**simplicial cohomology group** $\mathrm{SH}^n(R, G)$ of R with coefficients in G is the nth cohomology group of the complex obtained by applying the functor $\mathrm{Hom}_{\mathbb{Z}}(-, G)$ on the chain complex $(\mathrm{SC}_\bullet(R), d_\bullet)$.

Observe that the above chain complexes depend essentially on the way $\mathbb{B}$ is related to I. Hence, as for fundamental groups (see Section 4.2), different presentations of R may lead to different simplicial (co)homology groups. See [**5**, (5.2)] for an example.

If R is oriented by e, e'_1 and e'_2 with $\mathbb{B}_1$ and $\mathbb{B}_2$ semi-normed bases of A_1 and A_2, respectively, then $\mathbb{B}_R = \mathbb{B}_1 \cup \mathbb{B}_2$ and $\mathbb{B}_C = \mathbb{B}_1 \cap \mathbb{B}_2$ are semi-normed bases of R and C, respectively. This leads to the following results.

Proposition 4.1.1. *Let R be oriented by e, e'_1 and e'_2. Moreover, assume that A_1 and A_2 admit semi-normed basis. Then there exists a long exact sequence of simplicial homology groups:*

$$\cdots \longrightarrow \mathrm{SH}_n(C) \longrightarrow \mathrm{SH}_n(A_1) \oplus \mathrm{SH}_n(A_2) \longrightarrow \mathrm{SH}_n(R) \longrightarrow \mathrm{SH}_{n-1}(C) \longrightarrow \cdots$$

Proof. By the above discussion, R and C admit semi-normed bases. For $i \geq 0$, define

$$\begin{aligned} p_i : \qquad \mathrm{SC}_i(A_1) \oplus \mathrm{SC}_i(A_2) &\longrightarrow \mathrm{SC}_i(R) \\ ((\beta_1, \dots, \beta_i), (\gamma_1, \dots, \gamma_i)) &\longmapsto (\beta_1, \dots, \beta_i) - (\gamma_1, \dots, \gamma_i) \end{aligned}$$

The map $p_\bullet$ is a morphism of complexes. Moreover, let $\alpha = (\alpha_1, \dots, \alpha_i)$ be a generator of $\mathrm{SC}_i(R)$. Since $e'_1 R e'_2 = e'_2 R e'_1 = 0$, then, for each j, we have $\alpha_j \in \mathbb{B}_1$ or $\alpha_j \in \mathbb{B}_2$. Consequently, $\alpha \in \mathrm{SC}_i(A_1)$ or $\alpha \in \mathrm{SC}_i(A_2)$, whence p_i is surjective. Moreover, the kernel of p_i is exactly $\mathrm{SC}_i(C)$. We obtain a short exact sequence of complexes

$$0 \longrightarrow \mathrm{SC}_\bullet(C) \longrightarrow \mathrm{SC}_\bullet(A_1) \oplus \mathrm{SC}_\bullet(A_2) \longrightarrow \mathrm{SC}_\bullet(R) \longrightarrow 0 \tag{4.1}$$

from which we deduce the result. □

Proposition 4.1.2. *Let R be oriented by e, e'_1 and e'_2. Moreover, assume that A_1 and A_2 admit semi-normed basis. Then, for each abelian group G, there exists a long exact sequence of simplicial cohomology groups:*

$$\cdots \to \mathrm{SH}^n(R, G) \to \mathrm{SH}^n(A_1, G) \oplus \mathrm{SH}^n(A_2, G) \to \mathrm{SH}^n(C, G) \to \mathrm{SH}^{n+1}(R, G) \to \cdots$$

Proof. Since $\mathrm{SC}_i(R)$ is a free abelian group for all i, the sequence (4.1) splits and so the short sequence of complexes obtained by applying $\mathrm{Hom}_{\mathbb{Z}}(-, G)$ to the sequence (4.1) is exact. $\square$

4.2. Fundamental group. Let (Q, I) be a bound quiver. Recall from [**22**] that, given two vertices $x, y \in Q_0$, a relation $\rho = \sum_{i=1}^{n} \lambda_i \omega_i \in I \cap kQ(x, y)$ is called **minimal** provided $n \geq 2$ and $\sum_{i \in J} \lambda_i \omega_i \notin I \cap kQ(x, y)$ for every proper subset J of $\{1, 2, \ldots, n\}$. Let Q'_1 be the union of Q_1 with the set of inverse arrows $\alpha^{-1} : y \to x$ for each arrow $\alpha : x \to y$ in Q_1, a **walk** in (Q_0, Q_1) is a path in (Q_0, Q'_1). The **homotopy relation** in the set of walks on Q is the smallest equivalence relation such that:

(a) For each arrow $\alpha : x \to y \in Q_1$, $\alpha\alpha^{-1} \sim \epsilon_x$ and $\alpha^{-1}\alpha \sim \epsilon_y$;
(b) If $\sum_{i=1}^{n} \lambda_i \omega_i$ is a minimal relation, then $\omega_j \sim \omega_k$ for all $1 \leq j, k \leq n$;
(c) If u, v, w, w' are walks such that $w \sim w'$, then $uwv \sim uw'v$, whenever these compositions are defined.

A minimal relation ρ is **fundamental** if there does not exist a nontrivial path u and a minimal relation ω such that $\rho = \omega u$ or $\rho = u\omega$. One can see that the above homotopy relation does not change if we replace the condition (b) by

(b') If $\sum_{i=1}^{n} \lambda_i \omega_i$ is a fundamental minimal relation, then $\omega_j \sim \omega_k$ for all $1 \leq j, k \leq n$.

When dealing with homotopy, it is thus sufficient to consider only the fundamental minimal relations. Here, we give examples of fundamental minimal relations.

Example 4.2.1. Let Q be the quiver

$$\begin{tikzcd}
 & 2 \arrow[dl, "\alpha_1"'] & & & 6 \arrow[dl, "\delta_1"'] & \\
1 & & 4 \arrow[ul, "\beta_1"'] \arrow[dl, "\beta_2"] & 5 \arrow[l, "\gamma"'] & & 8 \arrow[ul, "\varepsilon_1"'] \arrow[dl, "\varepsilon_2"] \\
 & 3 \arrow[ul, "\alpha_2"] & & & 7 \arrow[ul, "\delta_2"] &
\end{tikzcd}$$

bound by the relations $\beta_1\alpha_1 - \beta_2\alpha_2 = \varepsilon_1\delta_1 - \varepsilon_2\delta_2 = \delta_1\gamma\beta_1 = \delta_2\gamma\beta_2 = 0$. The minimal fundamental relations are $\beta_1\alpha_1 - \beta_2\alpha_2$ and $\varepsilon_1\delta_1 - \varepsilon_2\delta_2$, while the relations $\gamma(\beta_1\alpha_1 - \beta_2\alpha_2)$ and $(\varepsilon_1\delta_1 - \varepsilon_2\delta_2)\gamma$ are minimal but not fundamental.

For a fixed vertex $x \in Q_0$, let $\pi_1(Q, x)$ be the fundamental group of the underlying graph of Q at the vertex x, that is the set of homotopy classes of closed walks on Q starting and ending at x. It is well-known (see [**25**], for instance) that $\pi_1(Q, x)$ is isomorphic to the free multiplicative nonabelian group on $\chi(Q) = |Q_1| - |Q_0| + 1$ generators. Now, let $N(Q, I, x)$ be the normal subgroup of $\pi_1(Q, x)$ generated by the elements of the form $wuv^{-1}w^{-1}$ where w is a path starting at x and u, v are two homotopic paths. The **fundamental group at** x of the bound quiver (Q, I) is defined to be $\pi_1(Q, I, x) = \pi_1(Q)/N(Q, I, x)$. If Q is connected, this definition does not depend on the choice of x. Accordingly, we write $\pi_1(Q, I) = \pi_1(Q)/N(Q, I)$, and call it the **fundamental group** of the bound quiver (Q, I). Similarly, we write $\pi_1(Q)$ instead of $\pi_1(Q, x)$.

On other hand, following [**5**], to a bound quiver (Q, I), we associate a CW complex $\mathcal{B}(Q, I)$ (called its **classifying space**) in the following way: the 0-cells are given by Q_0 and the n-cells are given by n-tuples $(\overline{\sigma_1}, \ldots, \overline{\sigma_n})$ of homotopy classes of nontrivial paths of (Q, I) such that the composition $\sigma_1\sigma_2 \cdots \sigma_n$ is a path of (Q, I)

not in I. Then, by [**5**, (3.3)], there is an isomorphism of groups

$$\pi_1(Q,I) \simeq \pi_1(\mathcal{B}(Q,I)) \tag{4.2}$$

We shall use this fact, together with Van Kampen's theorem, in the following proof.

PROPOSITION 4.2.2. *Let $R = kQ_R/I_R$ be a bound quiver algebra, oriented by three idempotents e, e'_1 and e'_2. Moreover, assume that $C = kQ_C/I_C$, $A_1 = kQ_{A_1}/I_1$ and $A_2 = kQ_{A_2}/I_2$ and that, for each fundamental minimal relation ω on (Q_R, I_R), all the paths occurring in ω are either contained in Q_C or contain no arrows from Q_C. Let $Q_C = \coprod_{j=1}^m Q_{C_j}$ with each Q_{C_j} connected, and set $I_{C_j} = I_R \cap kQ_{C_j}$. Then,*

$$\pi_1(Q_R, I_R) \simeq \frac{\pi_1(Q_{A_1}, I_{A_1}) \coprod \pi_1(Q_{A_2}, I_{A_2}) \coprod L_{m-1}}{\coprod_{j=1}^m \pi_1(Q_{C_j}, I_{C_j})}$$

where $\coprod$ denotes the free product of groups and L_{m-1} is the free group in $m-1$ generators.

PROOF. Let n_1, n_2, n_R, n and n_j be the number of vertices in $Q_{A_1}, Q_{A_2}, Q_R, Q_C$ and Q_{C_j} respectively. In particular, $n_R = n_1 + n_2 - n$. Let T_{C_j} be a maximal tree in Q_{C_j}, T_C be the disjoint union of the T_{C_j} and for $i \in \{1,2\}$, let T_i be a maximal tree in Q_{A_i} such that $T_1 \cap T_2 = T_C$. Such maximal trees always exist. In addition, $|(T_C)_1| = \sum_{j=1}^m |(T_{C_j})_1| = \sum_{j=1}^m (n_j - 1) = n - m$. For $i = 1, 2$, set

$$\hat{Q}_{A_i} = Q_{A_i} \coprod_{T_C} T_k \text{ with } k \neq i \text{ and } \hat{I}_i = I_R \cap k\hat{Q}_{A_i}.$$

Observe that $Q_R = \hat{Q}_{A_1} \coprod_{Q_C} \hat{Q}_{A_2}$. In addition, $|(\hat{Q}_{A_1})_0| = n_R$ and

$$|(\hat{Q}_{A_1})_1| = |(Q_{A_1})_1| + |(T_2)_1| - |(T_C)_1| = |(Q_{A_1})_1| + (n_2 - 1) - (n - m)$$

and a similar expression holds for $|(\hat{Q}_{A_2})_1|$. So, $\pi_1(\hat{Q}_{A_i})$ is the free group in $\chi(\hat{Q}_{A_i})$ generators, where $\chi(\hat{Q}_{A_i}) = \chi(Q_{A_i}) + (m-1)$. Then, $\pi_1(\hat{Q}_{A_i}) \simeq \pi_1(Q_{A_i}) \coprod L_{m-1}$.

Moreover, by our assumption on the fundamental minimal relations, we have an inclusion of groups $\iota_i : \pi_1(\hat{Q}_{A_i}, \hat{I}_i) \to \pi_1(Q_R, I_R)$. Also, the restriction to (Q_{A_i}, I_i) of the homotopy relation on $(\hat{Q}_{A_i}, \hat{I}_i)$ coincides with the homotopy relation on (Q_{A_i}, I_i). Thus $\hat{I}_i = I_i + J_i$ for some monomial ideal J_i and

$$\begin{aligned}
\pi_1(\hat{Q}_{A_i}, \hat{I}_i) &\simeq \pi_1(\hat{Q}_{A_i})/N(\hat{Q}_{A_i}, \hat{I}_i) \\
&\simeq (\pi_1(Q_{A_i}) \textstyle\coprod L_{m-1})/N(\hat{Q}_{A_i}, \hat{I}_i) \\
&\simeq [\pi_1(Q_{A_i})/N(Q_{A_i}, I_i)] \textstyle\coprod L_{m-1} \\
&\simeq \pi_1(Q_{A_i}, I_i) \textstyle\coprod L_{m-1}.
\end{aligned}$$

Now set $Q = \hat{Q}_{A_1} \cap \hat{Q}_{A_2}$ and $I = kQ \cap I_R$. We have

$$\begin{aligned}
\chi(Q) &= (|(T_1)_1| + |(T_2)_1| - 2|(T_C)_1| + |(Q_C)_1|) - n_R + 1 \\
&= |(Q_C)_1| - (n - m) + (m - 1) \\
&= (\sum_{j=1}^m \chi(Q_{Cj})) + (m - 1).
\end{aligned}$$

and so $\pi_1(Q) = (\coprod_{j=1}^{m} \pi_1(Q_{C_j})) \coprod L_{m-1}$. Moreover, by our assumption on the fundamental minimal relations, we have an inclusion of groups $\kappa_i : \pi_1(Q, I) \to \pi_1(\hat{Q}_{A_i}, \hat{I}_i)$ and $I = I_C + J_I$ for some monomial ideal J_I. So, $\pi_1(Q, I) = L_{m-1} \coprod (\coprod_{j=1}^{m} \pi_1(Q_{C_j}, I_{C_j}))$.

The next step is to use Van Kampen's theorem for bound quivers. For the sake of this, let $\hat{\mathcal{B}}_1$, $\hat{\mathcal{B}}_2$, $\mathcal{B}_R$ and $\mathcal{B}$ be the classifying spaces of $(\hat{Q}_{A_1}, \hat{I}_1)$, $(\hat{Q}_{A_2}, \hat{I}_2)$, (Q_R, I_R), (Q, I), respectively. Then, $\mathcal{B}_R = \hat{\mathcal{B}}_1 \cup \hat{\mathcal{B}}_2$ and $\mathcal{B} = \hat{\mathcal{B}}_1 \cap \hat{\mathcal{B}}_2$, where $\mathcal{B}$ is connected. Applying Van Kampen's theorem (see **[25]**) and the isomorphism of fundamental groups (4.2), we have

$$\begin{array}{rcl}
\pi_1(Q_R, I_R) & \simeq & \pi_1(\mathcal{B}_R) \\
& \simeq & \dfrac{\pi_1(\mathcal{B}_1) \coprod \pi_1(\mathcal{B}_2)}{\pi_1(\mathcal{B})} \\
& \simeq & \dfrac{\pi_1(\hat{Q}_{A_1}, \hat{I}_1) \coprod \pi_1(\hat{Q}_{A_2}, \hat{I}_2)}{\pi_1(Q, I)} \\
& \simeq & \dfrac{(\pi_1(Q_{A_1}, I_1) \coprod L_{m-1}) \coprod (\pi_1(Q_{A_2}, I_2) \coprod L_{m-1})}{(\coprod_{j=1}^{m} \pi_1(Q_{C_j}, I_{C_j})) \coprod L_{m-1}} \\
& \simeq & \dfrac{\pi_1(Q_{A_1}, I_1) \coprod \pi_1(Q_{A_2}, I_2) \coprod L_{m-1}}{\coprod_{j=1}^{m} \pi_1(Q_{C_j}, I_{C_j})}
\end{array}$$

□

We note that the main tool used in this proof is that the restriction of the homotopy relation on (Q_R, I_R) to (Q_C, I_C) coincides with the homotopy relation on (Q_C, I_C). This proof may thus clearly be generalized to this more general context.

The above formula allows to compute the fundamental group of several oriented algebras, but it does not include all oriented algebras. The following examples illustrate this point.

Examples 4.2.3.

(a) Let R be the algebra given by the bound quiver of Example 4.2.1. Let e, e'_1 and e'_2 be the sums of the primitive idempotents corresponding to the sets of vertices $\{4, 5\}$, $\{1, 2, 3\}$ and $\{6, 7, 8\}$ respectively. Clearly R is oriented by e, e'_1 and e'_2. Moreover, R satisfies to the conditions of Proposition 4.2.2. Since straightforward computations show that $\pi_1(Q_C, I_C)$, $\pi_1(Q_{A_1}, I_1)$ and $\pi_1(Q_{A_2}, I_2)$ are trivial, so is $\pi_1(Q_R, I_R)$ by Proposition 4.2.2.

(b) Let R be the algebra given by the quiver

$$\begin{tikzcd}
1 \arrow[r, "\alpha_1"] & 3 \arrow[d, shift right, "\beta"'] \arrow[d, shift left, "\gamma"] & 2 \arrow[l, "\alpha_2"'] \\
& 4 &
\end{tikzcd}$$

bound by the relations $\alpha_i(\beta - \gamma)$, for $i = 1, 2$. Let e, e'_1 and e'_2 be the sums of the primitive idempotents corresponding to the sets of vertices $\{3, 4\}$, $\{1\}$ and $\{2\}$ respectively. Clearly R is oriented by e, e'_1 and e'_2. However, R does not satisfy the conditions of Proposition 4.2.2 because the fundamental minimal relations $\alpha_i(\beta - \gamma)$ contain arrows from A'_i and C. Moreover, $\pi_1(Q_R, I_R)$, $\pi_1(Q_{A_1}, I_1)$ and $\pi_1(Q_{A_2}, I_2)$ are trivial, while $\pi_1(Q_C, I_C) \simeq \mathbb{Z}$. So the formula of Proposition 4.2.2 does not hold.

We end this section with an application of the above result to the computation of the first Hochschild cohomology group for some schurian oriented algebras (see Examples 2.4.1 (a)). Recall that if $A = kQ/I$ is a schurian algebra, then $\mathrm{H}^1(A) \simeq \mathrm{Hom}(\pi_1(Q,I), k^+)$ by [**10, 1**], where k^+ is the (abelian) additive group of k and the homomorphisms are group homomorphisms. This leads to the following result:

Proposition 4.2.4. *Assume R is oriented by e, e'_1 and e'_2 and satisfies to the conditions of Proposition 4.2.2. If R is schurian and $\coprod_{i=1}^{m} \pi_1(Q_{C_i}, I_{C_i})$ is trivial, then* $\mathrm{H}^1(R) \simeq \mathrm{H}^1(A_1) \oplus \mathrm{H}^1(A_2) \oplus k^{m-1}$.

Proof. Since R is schurian, then so are A_1, A_2 and C. Therefore

$$\begin{aligned}
\mathrm{H}^1(R) &\simeq \mathrm{Hom}(\pi_1(Q_R, I_R), k^+) \\
&\simeq \mathrm{Hom}(\pi_1(Q_{A_1}, I_{A_1}) \amalg \pi_1(Q_{A_2}, I_{A_2}) \amalg L_{m-1}, k^+) \\
&\simeq \mathrm{Hom}(\pi_1(Q_{A_1}, I_1), k^+) \oplus \mathrm{Hom}(\pi_1(Q_{A_2}, I_2), k^+) \oplus k^{m-1} \\
&\simeq \mathrm{H}^1(A_1) \oplus \mathrm{H}^1(A_2) \oplus k^{m-1}
\end{aligned}$$

□

Example 4.2.5. Let R be the algebra of Examples 1.2.1 (b). Theorem 2.3.2 cannot be used to compute the Hochschild cohomology groups of R. However, R is schurian and clearly satisfies (1.1) and the conditions of Proposition 4.2.2 since C is semisimple. Thus $\mathrm{H}^1(R) \simeq \mathrm{H}^1(A_1) \oplus \mathrm{H}^1(A_2) \oplus k \simeq k$ by Proposition 4.2.4.

ACKNOWLEDGEMENTS. This paper was started while the first author visited the Universities of Sherbrooke and Bishop's under a post-doctoral fellowship, and while the second and the third author respectively had an ISM and NSERC graduate studies fellowship; unfortunately, the write-up of the results has been delayed for quite a while. The authors warmly thank Ibrahim Assem for his precious help and patience.

References

[1] I. Assem and J. A. de la Peña, *The fundamental groups of a triangular algebra*, Comm. Algebra **24** (1996), no. 1, 187–208.

[2] B. Bendiffalah and D. Guin, *Cohomologie de l'algèbre triangulaire et applications*, J. Algebra **282** (2004), no. 2, 513–537.

[3] D. J. Benson, *Representations and cohomology. II: Cohomology of groups and modules*, Cambridge Studies in Advanced Mathematics, no. 31, Cambridge University Press, Cambridge, 1998.

[4] K. Bongartz and P. Gabriel, *Covering spaces in representation theory*, Invent. Math **65** (1981-1982), no. 3, 331–378.

[5] J. C. Bustamante, *The classifying space of a bound quiver*, J. Algebra **277** (2004), no. 2, 431–455.

[6] H. Cartan and S. Eilenberg, *Homological algebra*, Princeton mathematical series, no. 19, Princeton University Press, Princeton, 1956.

[7] C. Cibils, *Cohomology of incidence algebras and simplicial complexes*, J. Pure Appl. Algebra **56** (1989), 221–232.

[8] ______, *Hochschild cohomology algebra of radical square zero algebras*, Algebras and Modules II (Providence, RI), CMS Conf. Proc., no. 24, Amer. Math. Soc., 1998, pp. 93–101.

[9] ______, *Tensor Hochschild homology and cohomology.*, Interactions between ring theory and representations of algebras (Murcia) (New-York), Lecture Notes in Pure and Appl. Math., vol. 210, Dekker, 2000, pp. 35–51.

[10] J. A. de la Peña and M. Saorín, *The first Hochschild cohomology group of an algebra*, Manuscripta Math. **104** (2001), 1–12.

[11] S. Gastaminza, J.A. de la Peña, M.I. Platzeck, M.J. Redondo, and S. Trepode, *Algebras with vanishing Hochschild cohomology*, J. Algebra **212** (1999), no. 1, 1–16.
[12] M. Gerstenhaber, *The Cohomology structure on an associative ring*, Ann. of Math. **78** (1963), no. 2, 267–288.
[13] E. Green and Ø. Solberg, *Hochschild cohomology rings and triangular rings, Representations of algebra*, Beijing Norm. Univ. Press **I** (2002), 192–200.
[14] E. L. Green, E. N. Marcos, and N. Snashall, *The Hochschild cohomology ring of a one point extension.*, Comm. Algebra **31** (2003), no. 1, 357–379.
[15] D. Happel, *Triangulated Categories in the Representation Theory of Finite-Dimensional Algebras*, London Mathematical Society Lecture Note Series, 119, Cambridge University Press, Cambridge-New York, 1988.
[16] D. Happel, *Hochschild cohomology of finite-dimensional algebras*, Lecture Notes in Mathematics, no. 1404, pp. 108–126, Springer-Verlag, Berlin Heidelberg New York Tokyo, 1989.
[17] K. Igusa, M. I. Platzeck, G. Todorov, and D. Zacharia, *Auslander algebras of finite representation type*, Comm. Algebra **13** (1987), no. 1-2, 377–424.
[18] J. Lévesque, *Nakayama oriented pullbacks and stably hereditary algebras*, J. Pure Appl. Algebra **212** (2008), no. 5, 1149-1161.
[19] S. Liu and J. P. Morin, *The strong no loop conjecture for special biserial algebras*, Proc. Amer. Math. Soc. **132** (2004), no. 12, 3513–3523.
[20] J.-L. Loday, *Cyclic homology*, Grundlehren der Mathematischen Wissenschaften, no. 301, Springer-Verlag, Berlin, 1992.
[21] Ma. I. R. Martins and J. A. de la Peña, *Comparing the simplicial and the Hochschild cohomologies of a finite dimensional algebra*, J. Pure Appl. Algebra **138** (1999), no. 1, 45–58.
[22] R. Martínez-Villa and J. A. de la Peña, *The universal cover of a quiver with relations*, J. Pure Appl. Algebra **30** (1983), no. 3, 873–887.
[23] S. Michelena and M. I. Platzeck, *Hochschild cohomology of triangular matrix algebras*, J. Algebra **233** (2000), no. 2, 502–525.
[24] M. J. Redondo, *Hochschild cohomology: some methods for computations*, Resenhas IME-USP **5** (2001), no. 2, 113–137.
[25] J.J. Rotman, *An introduction to Algebraic topology.*, Graduate Texts in Mathematics, no. 119, Springer-Verlag, New York Berlin Heidelberg, 1988.
[26] A. Skowroński, *Simply connected algebras and Hochschild cohomologies.* Proceedings of the Sixth International Conference on Representations of Algebras (Ottawa, ON, 1992), 18 pp., Carleton-Ottawa Math. Lecture Note Ser., 14, Carleton Univ., Ottawa, ON, 1992.

J. C. Bustamante; Departamento de Matemáticas, Universidad San Francisco de Quito, Quito, Ecuador.
E-mail address: `juanb@usfq.edu.ec`

J. Dionne; Département de Mathématiques, Université de Sherbrooke, Sherbrooke, J1K 2R1, Québec, Canada.
E-mail address: `julie.dionne2@usherbrooke.ca`

D. Smith; Department of Mathematics, Bishop's University, Sherbrooke, Québec, Canada, J1M0C8.
E-mail address: `unracinois@hotmail.com`